



Hunt Institute for Botanical Documentation  
5th Floor, Hunt Library  
Carnegie Mellon University  
4909 Frew Street  
Pittsburgh, PA 15213-3890  
Telephone: 412-268-2434  
Email: [huntinst@andrew.cmu.edu](mailto:huntinst@andrew.cmu.edu)  
Web site: [www.huntbotanical.org](http://www.huntbotanical.org)

The Hunt Institute is committed to making its collections accessible for research. We are pleased to offer this digitized item.

*Usage guidelines*

We have provided this low-resolution, digitized version for research purposes. To inquire about publishing any images from this item, please contact the Institute.

*About the Institute*

The Hunt Institute for Botanical Documentation, a research division of Carnegie Mellon University, specializes in the history of botany and all aspects of plant science and serves the international scientific community through research and documentation. To this end, the Institute acquires and maintains authoritative collections of books, plant images, manuscripts, portraits and data files, and provides publications and other modes of information service. The Institute meets the reference needs of botanists, biologists, historians, conservationists, librarians, bibliographers and the public at large, especially those concerned with any aspect of the North American flora.

Hunt Institute was dedicated in 1961 as the Rachel McMasters Miller Hunt Botanical Library, an international center for bibliographical research and service in the interests of botany and horticulture, as well as a center for the study of all aspects of the history of the plant sciences. By 1971 the Library's activities had so diversified that the name was changed to Hunt Institute for Botanical Documentation. Growth in collections and research projects led to the establishment of four programmatic departments: Archives, Art, Bibliography and the Library.

— Cladivum. Brown p. 114. *of adoni* (synonym).  
Scipi (rec. waterlily) *pseudo cyathum* pl. 7. et contra ferunt. non adonis. <sup>39</sup> quaternary sand,  
cal. aquina 5 valv 1 flor. Cor. Sic ampliatus fls d Brown. C. quinque 6. piece interrate  
con. 1. valv 1. apothecium ovarianum. bellarianum atro 2. interna dors. corolla ling. 1 florae, quae 5 infruct. abortant. sed stig 2 habet  
filamenta stativa, quae brown except pro nectando.  
stem 2. var 1. lilyt. 1. stigmat 3 cyl.  
Semp. ovat. *fragi*

# TRIANDRIA DIGYNIA.

54. SCIRPUS\*. Mich. Tournef. 300. Scirpociperus. caulis. 3gney.  
caulis rotund. emarginati. praerootatis cibida. Mich. 31. 38. pira spica.  
Scleria. mich. Tab 31. 20pt. a furo sumis. abz. v. 10. 7

CAL. *Spica imbricata*: squamis ovatis, plano-inflexis, flores distincti  
gumentibus.

**Cor.** nulla.

STAM. Filamenta tria, tandem longiora. Antheræ oblongæ.

Pist. Germen minimum. Stylus filiformis, longus. Stigmata tria,  
capillaria.

P E R. nullum.

*S E M. unicum , triquetrum , acuminatum , villis calyce brevioribus  
instructum. en inde at variat de pictu-*

**O**b s. villi isti seminis in aliis ad apicem, in aliis ad basin semini annectuntur.

55. ~~ERIOPHORUM~~ \* Linagrostis <sup>c.B. Tab. c.B.</sup> Mich. 31. Tournef. Cormaralle <sup>multivirg.</sup>  
*Pleurosticus* L. <sup>supreme-</sup>  
flores inspiciat tent. lund. <sup>depi</sup> vaginæ fol. integ. non lutea  
grandj. varia ubi sedet.  
b. i. s. l. i. s. i. s. i. s. i. s.

**C A L.** *Spica imbricata*; *squamis ovato-oblongis*, *plano-inflexis*, *flores* *distinguenteribus*.

**C**or. nulla.

**STAM.** *Filamenta tria*, capillaria. *Antheræ erectæ*, oblongæ.

Pist. Germen minimum. Stylus filiformis, longitudine squamæ calycis. Stigmata <sup>2 in mhd.</sup> tria, stylo longiora, reflexa.

P E R. nullum.

SEM. triquetrum, acuminatum, villis spica longioribus instru-  
tum. villi circa ovarium non concrecentia glandem nisi per cruentationem.

## II. DIGYNIA.

56. CORNUCOPIÆ t. Cornucopioides Scheuch. 3.

**CAL.** Perianthium commune, monophyllum, infundibuliforme, <sup>15.</sup> multiflorum  
maximum: ore crenato, obtuso, patenti-erecto.

*Gluma uniflora*, bivalvis: *valvulis oblongis*, obtuse acuminatis, <sup>water nitre  
Dioscorea</sup> æqualibus.

**C**or. univalvis, figurā, magnitudine & situ valvulae calycis simil-  
lima.

**S T A M.** *Filamenta tria*, capillaria. *Atheræ oblongæ*,  
*P. G. turbinatum*. *Stuli duo*, capillares. *Stigmata cirrosa*.

P I S T. Germen turbinatum. Styli duo, & P. all. C. all. sumen includens

P E R. nullum. Corolla semen includens.  
S E M. unicum turbipatum hinc convexum, inde planum.

Nazia. Adams. green <sup>10</sup> spicat lvs. set. achen. Convolv. Linn.  
Car. pilos. fl. spic. hemph. Cal. green 2 valv. ovat. capill. awl. Cor. 2 valv. awl. ang.  
ovate 3. styl. 2. awl.

22 TRIANDRIA DIGYNIA.

Vasus. Ddore ut heta indra 2 ad 3 spithamea - alit.

57. NARDUS \*. vide Scheuch. 2 : 10. Mont. 31.

C A L. nullus.

**COR.** bivalvis, *Valvula exterior* lanceolato-linearis, longa, sensim in aristam desinens, ventre amplectens minorem: *V. minor* interior, linearis, in aristam breviorem desinens.

*Stam.* Filamenta tria, capillaria, corollâ breviora. Antheræ oblongæ.

P I S T. Germen oblongum. Styli duo, capillares, reflexi, pubescentes. Stigmata simplicia.

**P**E.R. Corolla adnascitur semini, nec dehiscit.

S E M. unicum, tectum, linearis-longum, utrinque acuminatum, superne angustius.

58. *calamagrostis*  
**SACCHARUM** †. CANNE A SUCRE *albus*

**C A L.** nullus, sed lanugo flori longior, florem unicum includens.  
**C O R.** bivalvis: *Valvulae* oblongo-lanceolatae, acuminatae, erectae.

CAL. nullus, sed lanugo flori longior, florem unicum includens.  
COR. bivalvis: Kelazula oblonga longior.

*Cor. bivalvis: Valvulae oblongo-lanceolatae, acuminatae, erectae,  
concavae, aequales, muticae.*

STAM. *Filamenta tria, capillaria, longitudine corollæ. Antheræ oblongiusculæ.*

P I S T. Germen subulatum. Styli duo, cirrosi. Stigmata simplicia.  
P E R. nullum. Corolla foveat semen.

**S E M:** unicum oblongum angustum

*E. M. unicum, oblongum, angustum, acuminatum.*

9. PHALARIS \*. Phalaroidea Scheuch. 2 : 3. Mont. 44.

CAL. *Gluma uniflora*, bivalvis, maxima, compressa, obtusa: *valvulis* navicularibus, compressis, carinatis, superne obtusioribus, marginibus rectis, parallele conniventibus.

*Cor. bivalvis, calyce minor: Valvula exteriore oblonga, acuminate, convoluta; interiore minore.*

T A M. *Filamenta tria*, *capillaria*, *calyce breviora*. *Antheræ oblongæ.*

I S T. *Germen subrotundum. Styli duo, capillares. Stigmata villosa.*

E.R. Corolla circumnascitur semini crustinae — labifida


E. *Corolla circumfalcitum lemni crinitæ instar, nec dehiscit.*  
E. M. *unicum, tectum, glabrum, è rotunda utrinque cum in tum*

B S. in quibusdam speciebus, Gluma calycina minor, intra caly-  
cem descriptum, bivalvis, acuminata, minima, adest.

~~the genus~~  
O. ~~con. major~~ PLHEUM \* Tiphooidea Scheuch. 2 : 5. Staphylin.

C A L. *Gluma uniflora*, bivalvis, oblonga, linearis, compressa, apice bicuspidate dehiscens: *valvulis* rectis, concavis, compressis, amplexantibus.

*Cor. bivalvis, calyce brevior: Valvula exteriore amplexante inter-*



~~mucilosa~~: adans. geogr. 10. hin. gramen. mon. f. 3. l. 2. f. 10.  
~~mibra~~

### TRIANDRIA DIGYNIA

23

STAM. Filamenta tria, capillaria, calyce longiora. Antheræ oblongæ, bifurcatæ.

PIST. Germen subrotundum. Styli duo, capillares, reflexi. Stigmata plumosa.

PER. nullum. Calyx & corolla includentia semen.

SEM. unicum, subrotundum.

### 61. ALOPECURUS\*. Myosuroidea Scheuch. 2: 5. 6.

*lamea membr. brev. int.*

CAL. Gluma uniflora, bivalvis: Valvulis ovato-lanceolatis, concavis, compressis, æqualibus.

COR. univalvis: Valvula concava, longitudine calycis. Aristæ longa, dorso valvulae propè basin inserta.

STAM. Filamenta tria, capillaria. Antheræ utrinque bifurcæ.

PIST. Germen subrotundum. Styli duo, cirrofisi, reflexi, calyce longiores. Stigmata simplicia.

PER. Corolla includit semen.

SEM. unicum, subrotundum, tectum.

### 62. PANICUM\*. P<sup>an</sup>ANIS. Panicea Scheuch. 2: 2.

*Exupes, pugnac. Thymo. pedunc. dist.*

*Ledactylon. 1. Plan. corona membr. brev.*

CAL. Involucrum uniflorum, polypyllum: foliolis capillaribus, inæqualis insertionis.

Gluma uniflora, trivalvis: Valvulis ovato-acuminatis: tertia minima à tergo alterius posita.

COR. bivalvis: Valvulae ovato-acuminatæ, altera minor planior.

STAM. Filamenta tria, capillaria, brevia. Antheræ oblongæ.

PIST. Germen subrotundum. Styli duo, capillares. Stigmata. plumosa.

PER. Corolla adnascitur semini, nec dehiscit.

SEM. unicum, tectum, subrotundum, inde planiusculum.

OBS. Aristæ terminant in quibusdam corollæ valvulae planiorem.  
*Species datur involucro destituta.*

### 63. MILIUM\*. MILLER. Tournef. 298.

*Milium parvum. 1. florum. calix 2 v. alv. corona floris membrana magna prævara*

*De plan. libid. Hipp. Keph. Nostros. Hipp.*

CAL. Gluma uniflora, bivalvis: Valvulis ovatis, acuminatis.

COR. bivalvis, calyce minor: Valvulae ovatæ, altera earum minor.

STAM. Filamenta tria, capillaria, brevissima. Antheræ oblongæ.

PIST. Germen subrotundum. Styli duo, capillares. Stigmata penicilliformia.

PER. Corolla adnascitur semini, nec dehiscit.

SEM. unicum, tectum, subrotundum.

OBS. Milium calyce bivalvi à Panico calyce trivalvi differt.

*Milium vulg. unalb. id ac parvum. 2 florum. cor. pilos. brev.*

B iiiij

Melica uniflora. Inflorescens est inter omnia graminea gen. Valvulae oblongae ex 2 à 3 membranis levata, clavatice vel levigata, fimbriis debitis  
comite membranis triang. levata et tenui corona 2 pedicellis velatis.

### TRIANDRIA DIGYNIA.

24

65. DED. C. B. 26.

Melica Ital.  
ex numero  
longhi.

64.

### MELICA \*. Fl. Lapp §. 36.

CAL. Gluma uniflora, bivalvis: Valvulis ovatis, concavis, fere æqualibus.

COR. bivalvis: Valvulae ovatae, muticæ: altera concava: altera vero plana.

STAM. Filamenta tria, capillaria, longitudine floris. Antheræ oblongæ, utrinque bifurcatæ.

PIST. Germen ovato-turbinatum. Styli duo, setacei, patentes. Stigmata oblonga villosa.

PER nullum. Corolla includit semen, demittit.

SEM. unicum, ovatum.

OBS. Corpusculum pedunculatum intra calycem, juxta corollam (an floris rudimentum) characterem essentialiem præbet: hoc constat duobus flosculis truncatis, alternis: glumis convolutis, pellucidis.

Agrostis. d. 20. 65. AGROSTIS. vide Scheuch. 3: 11. 9.

agrostis sp. de  
semita aristata  
coron. membr.  
magna.

CAL. Gluma uniflora, bivalvis, acuminata.

COR. bivalvis, acuminata, vix longitudine calycis, altera major aristata.

STAM. Filamenta tria, capillaria, corollâ longiora. Antheræ furcatæ.

PIST. Germen subrotundum. Styli duo, reflexi, villosi. Stigmata consimilia.

PER. Corolla adnascitur semini, nec dehiscit.

SEM. unicum, subrotundum, utrinque acuminatum.

### — 66. LAGURUS †. Alopecuroidea Scheuch. 2: 4.

fl. spicata. Cor. membr. magna integ.

CAL. Gluma uniflora, bivalvis: Valvulis oblongis, linearibus patulis, tenuissimis, desinentibus singulis in setam villis pennatam.

COR. Gluma bivalvis, calyce crassior: Valvula exteriore longiore, terminata Aristis duabus, parvis, rectis; Aristâ tertiatâ è medio dorso valvula ejusdem reflexo-torta. Valvula interiore parva, acuminata.

STAM. Filamenta tria, capillaria. Antheræ oblongæ.

PIST. Germen turbinatum. Styli duo, setacei. Stigmata villosa.

PER. nullum. Corolla semini adnascitur.

SEM. solitarium, oblongum, tectum, aristatum.

67. DACTYLIS \*. Roy. Lugdh. 56. Nom. id. ac Sartorius  
quodammodo  
coron. membr. magno integ.

CAL. Gluma uniflora aut biflora subrotunda, obtusa, bivalvis, flosculis major: valvulis æqualibus, exteriore ventricosiore.

COR. Gluma obtusa, æqualis, bivalvis, calyce tecta. Valvula exteriore latiore ventricosiore; interiore planiore.

— 68. Oryza. a. day. AIRA  
oblonga. integrifolia.  
GAL. Gluma biflora, bivalvis æqualibus.  
COR. bivalvis: Valvula aristata flore vix duplo lata.  
STAM. Filamenta tria, oblongæ, utrinque furcatae.  
PIST. Germen ovatum, bescentia.  
PER. nullum. Corolla includit semen.  
SEM. subovatum, tectum.  
OBS. Datur species, in qua sub eodem calyce communis.

69. herba grana POA  
Cor. membr. integ.  
CAL. Gluma multiflora, cham ovato-oblongam  
COR. bivalvis: Valvula calyce paulo longiore  
STAM. Filamenta tria,  
PIST. Germen subrotundum, consimilia.  
PER. Corolla adnascitur  
SEM. unicum, oblongum, tectum.

70. BRIZA \*. Tremula  
Cor. membr. twining.  
CAL. Gluma multiflora, tam disticham colligens  
COR. bivalvis: Valvula perior minima, plana,  
STAM. Filamenta tria, c. PIST. Germen subrotundum, mata plumosa.  
PER. Corolla immutata  
SEM. unicum, subrotundum.

*Sestaria*: Scopoli Radix perenne fibra Plantae capitatae 1/2 p. in spiculatum  
culmi plur. / capitulo non ramosci aedeo, capitulo 2 ad 3 foliis exstans 1/2 pedis  
Spica parva terminalia, valgo capitis, ita varia. Corolla 2. Calyx 2 flor. 2 valv. siugulis aristatis.  
Con. 2 valv. valv. ante bin. major. 3 arista. dracaenae minima. calyx intus 2. 25 annos. ad Bay  
In auctor. Iridac. *hodie*

### TRIANDRIA DIGYNIA.

STAM. Filamenta tria, capillaria, longitudine corollæ. Antheræ aed. aristis bifurcæ.

PIST. Germen turbinatum. Styli duo, capillares, patentes. Stigmata plumosa.

PER. nullum. Calyx Corollaque semen includentes, dimitentes.

SEM. solitarium, subrotundum, hinc depresso, inde convexum, nudum.

OBS. Species alias calyce unifloro, aliæ bifloro sunt.

68. *Apnis adau. AIRA* \*. Fl. Lapp. §. 47. aera. dix. v. Loliu.

*abris Thys. avens vitium* *pratinus avens calyx copiellus*

CAL. Gluma biflora, bivalvis: Valvulis ovato-lanceolatis, acutis, æqualibus.

COR. bivalvis: Valvulae calyci simillimæ: altera sæpe exserente aristam flore vix duplo longiore.

STAM. Filamenta tria, capillaria, longitudine floris. Antheræ oblongæ, utrinque furcatæ.

PIST. Germen ovatum. Styli duo, setacei, patentes. Stigmata pubescentia.

PER. nullum. Corolla includens semen adnascitur.

SEM. subovatum, tectum.

OBS. Datur species, in qua etiam mares miscentur hermaphroditis sub eodem calyce communi; tum mas est aristatus, femina mutica.

69. *herba. grana. dr. in quo. tig. Thys. anaphi aegypt. apnis hippocast. cotyled. dax. rebal apis*  
*con. monb. his. eue* POA \*. vide Scheuch. 4: 17. *Paniculata heit.*

CAL. Gluma multiflora, bivalvis, mutica, flores in spicam disticham ovato-oblongam colligens: Valvulis ovato-acuminatis.

COR. bivalvis: Valvulae ovato-acuminatæ, concavæ, compressæ, calyce paulo longiores, muticæ.

STAM. Filamenta tria, capillaria. Antheræ bifurcatæ.

PIST. Germen subrotundum. Styli duo, reflexi, villosi. Stigmata consimilia.

PER. Corolla adnascitur semini, nec dehiscit.

SEM. unicum, oblongum, acuminatum utrinque, compressum, tectum.

70. BRIZA \*. Tremula Scheuch. 4. 7. 8. 9. 10. Phalaroidis *in quo*  
*sp. Mont. 34. 39. Tremula heit. ac foas.*  
*con. in ibi. tunc.*

CAL. Gluma multiflora, bivalvis, patens, flores in spicam cordatam disticham colligens: valvulis cordatis, concavis, æqualibus.

COR. bivalvis: Valvula inferior magnitudine & figura calycis; superior minima, plana, subrotunda, ventrem alterius claudens.

STAM. Filamenta tria, capillaria. Antheræ oblongæ.

PIST. Germen subrotundum. Styli duo, capillares, recurvi. Stigmata plumosa.

PER. Corolla immutata continet semen, dehiscit, demittit.

SEM. unicum, subrotundum, compressum, minimum.

W genus  
ac. Boe  
er me.

71.

71. UNIOLA f. vide Pluk. 32. f. 6.  
*car. membr. med.*

Car. membr. Med.

CAL. *Gluma* multiflora, sexvalvis: *Valvulis* distiche imbricatis, fere subulatis, compressis, navicularibus, carinulatis, altera alteram claudente: ultimo pari multifloro: continente spicam ovatam, compresso-planam, margine acutam.

**C**or. *Gluma bivalvis*: *Valvulae lanceo lato-compressæ*, *calycinis similes*: *interiore valvula parum supereminente.*

STRAM. *Filamenta tria, capillaria. Antheræ oblongæ, lineares.*

Pist. Germen conicum. Styli duo, errecti, simplices. Stigmata pubescentia.

PER. nullum. *Corolla* includit semen.

### S E M. unicum, ovato-oblongum.

72. CYNOSURUS \*. Cristata. Scheuch. 2. 1. vid.  
*Can. Marbr. med.* Mont. 23.

Mont. 23.

CAL. *Involucrum* partiale, laterale, triphyllum, magnum: foliolo  
pennato, persistente.

*Gluma multiflora*, bivalvis : *valvulis linearibus*, acuminatis, æqualibus.

**COR.** bivalvis: exterior concava, longior, aristata; interior mutica.

**S T A M.** Filamenta tria, capillaria. Antheræ oblongæ.  
**P I S T.** Germen turbinatum. Styli duo, villosi, reflexi. Stigmata

P E R. nullum. Corolla arcte obvolvens semen nec debiscens.

**S E M.** unicum, oblongum, utrinque acuminatum

~~w. gerry~~ 73. BROMUS\*. Mont. 32. Festuca Scheuch. 5: 2. 10. 11. <sup>spicules. Disk. are flatish awl-shaped, so called awl awns.</sup>  
~~as festuca~~ <sup>aveneria-huic</sup> 2. Aegilops Dill. gen. 3.  
~~animes~~ <sup>corona sub dividata.</sup>

CAL. *Gluma* multiflora, bivalvis, patens, flores in spicam ovato-oblongam colligens; *valvulis* ovato-oblongis, acuminatis, multicisi, inferiore minore.

**C**OR. bivalvis: *Valvula inferior* major, magnitudine & figura ca-  
lycis, concava, obtusa, aristam infra apicem rectam emittens,  
supra aristam bifida *V. superior* lanceolata, parva, mutica.  
**S**TAM. Filamenta tri-

STAM. Filamenta tria, capillaria, corollâ breviora. Antheræ oblongæ.  
PIST. Germen turbinatum. Styli duo, breves, reflexi, villosi. Stigmata simplicia.

P E R. Corolla arctissime clausa adnascitur nec debifit

S E M. unicum, oblongum, tectum, hinc convexum, inde fulcatum.

91. Elymus. fl. spirali. ex membr. ex 2 involucr. lateris  
et nonnum. cal. involuc. 2 phyllo. spiralis germinis. cal. prope ②  
Panicea. con. 2 ser. ap. ext. lato. -  
Miki eti. vites petes. 6-10 cm. altos. Reticuli 1. r. exalt. reticuli. Thresher.

TRIANDRIA DIGYNIA.

27

*Lat. Mor. f. 8. 2. 7. f. 11. 16.*

74. FESTUCA \*. Dill. gen. 5. Bromoides Scheuch. 5: 13-18.

CAL. *Gluma* multiflora, bivalvis, erecta, flores in spicam tenuem continens: *valvulis* linearis-oblongis, acuminatis, muticis, inferiore minore.

COR. bivalvis: *Valvula* inferior major, figurâ calycis, magnitudine calycem superans, teretiuscula, acuminata, in Aristam rectam desinens.

STAM. *Filamenta* tria, capillaria, corollâ breviore. *Antheræ* oblongæ.

PIST. *Germen* turbinatum. *Styli* duo, breves, reflexi. *Stigmata* simplicia.

PER. *Corolla* arctissime clausa adnascitur, nec dehiscit.

SEM. unicum, tenui-oblongum, utrinque acutissimum, longitudinaliter sulca notatum.

OBS. antecedenti valde affinis est; aristâ terminatrice distincta.

75. AVENA \*. AVOINE. Tournef. 267. Malp. 138. *Avenacea* Scheuch. 4: 15-22. 5: 1. Dill. gen. 2.

CAL. *Gluma* multiflora, bivalvis, flores laxos colligens: *valvulis* lanceolatis, acutis, ventricosis, laxis, magnis, muticis.

COR. bivalvis: *Valvula* inferior calyce durior, calycis magnitudine, teretiuscula, ventricosa, utrinque acuminata, è dorso Aristam spiraliter intortam, geniculo veluti reflexam emittens.

*Nectaria* duo, ovato-lanceolata, à latere superiore germini insidentia.

STAM. *Filamenta* tria, capillaria. *Antheræ* oblongæ, bifurcatæ.

PIST. *Germen* obtusum. *Styli* duo, reflexi, pilosi. *Stigmata* simplicia.

PER. *Corolla* arctissime clausa adnascitur, nec dehiscit.

SEM. unicum, tenui-oblongum, utrinque acuminatissimum, sulco longitudinali notatum.

OBS. Flores intra communem calycem in quibusdam duo, in aliis plures. Arista è dorso corollæ articulata, intorta, constituit, essentialm characterem. Datur species calyce uniglumi, arista terminatrice.

76. ARUNDO \*. CANNE ou ROSEAU. Scheuch. 3: 14. 3.

CAL. *Gluma* uni-vel multi-flora, bivalvis, erecta: *Valvulis* oblongis, acuminatis, muticis; alterâ breviore.

COR. bivalvis: *Valvulae* longitudine calycis, oblongæ, acuminatae, è quarum basi lanugo longitudine fere floris assurgit.

STAM. *Filamenta* tria, capillaria. *Antheræ* utrinque bifurcatæ.

PIST. *Germen* oblongum. *Styli* duo, capillares, reflexi, villosi. *Stigmata* simplicia.

PER. *Corolla* adnascitur semini, nec dehiscit.

SEM. unicum, oblongum, utrinque acuminatum, basi pappo longo instructa.

Bambu. Maumba. My. h. na. arundo canadensis. ex radice altitudine 50 m. fl. graminea vaginata. aperte. flor. panicul. grossul. rotund. arundinarii. cum 3 clavatibus.



80. HORDEUM \*. ORGE. Tournef. 295. Scheuch. I: 3. *spicig. Rhynchon.*  
*lun. membr. ond. cum 2 aundis. latibus*

CAL. *Involucrum partiale, triflorum, hexaphyllum, foliolis linearibus, acuminatis, erectis, duobus singulo flori subjectis. Gluma nulla.*

COR. bivalvis: *Valvula inferior ventricosa, angulata, ovato-acuminata, calyce longior, desinens in Aristam longissimam: V. interior lanceolata, plana, minor.*

STAM. *Filamenta tria, capillaria, corollâ breviora. Antheræ oblongæ.*

PIST. *Germen ovato-turbinatum. Styli duo, villosi, reflexi. Stigmata similia.*

PER. *Corolla circumnascitur semini, nec dehiscit.*

SEM. *oblongum, ventricosum, angulatum, utrinque acuminatum, inde sulca longitudinali notatum.*

OBS. *In quibusdam omnes tres flores, involucro contenti, hermaphroditi sunt fertiles; in aliis flores laterales mares, sunt intermedius autem solus fertilis hermaphroditus. Hinc H. distichon, & Gramina bordeacea veterum.*

### III. TRIGYNA.

81. ERIOCALLON\*. Gron. Virg. I 4. *ad. staticum*  
*folia raddi statice vel graminis in 2 spiculis. in una spica petio fl. majora*  
*anaplecta haeroph. folia fere tripl. pera quamquidam fructus super basim cauli pta.*

CAL. *Perianthium commune, globoso-depressum, imbricatum squamis lanceolatis æqualibus persistentibus.*

COR. *Universalis uniformis convexa.* *en hinc ovario insidet.*  
*Propria tripetala: Petalis lanceolatis, obtusis, apice, villosis, basi attenuatis connatisque in pedicellum styliforme pilosum.*

STAM. *Filamenta tria, capillaria, germini infistentia. Antheræ oblongæ versatiles.*

PIST. *Germen tenue intra corollam, sub stamina. Styli tres capillares breves. Stigmata simplicia.*

PER. *nullum. Calyx immutatus.*

SEM. *solitaria, corollâ volitantia.*

REC. *Paleæ magnitudine & figura squamarum calycis, unifloræ, plurimæ.*

Polyg 89. apluda. Linn. auct. *Thunberg en die id ac Zeydy Brown*  
*mention. cal. gluma 3 flora 2 valv. pro parte flore 1. femina sibili et 2. male pedicello connex*  
*floribus cal. o.*  
*con gluma 2 valv. exar. orat mutata. In Schizanthus cordata*  
*exar. orat. styl. 1. pubesc. tig. 1. styl. in lobi. 1 styl.*  
*Sanct. 1. orat. long.*  
*male 2. in eod. pedicello.*  
*cal. o.*  
*con. 2 valv. exar. mutata*  
*styl. 3.*  
*flo. sensu vaginam attenuante ingrediendum*  
*pannula pera angusta, laterali.*



~~—Sectaria 24. domingensis Jaeg. Cal. 4 dent. Cor. hypocrateus. Stig. I. Drupe. Nua 2 locul.  
—Aquatica 24. aquatica Jaeg. Cal. 4 dent. Cor. rotata 4 f. Nua 1 locul. polyp.~~

TETRANDRIA MONOGYNIA. 31

~~Chamaea Jaeg. 24. spinosa. Cal. superus 4 pd. Cor. tubularis 4 pd. Stam. 4. Drupe  
nua 2 locul.~~

Classis IV.

# TETRANDRIA.

## I. MONOGYNIA.

S. Blatt.

Lepidocarpodendron ad

—85. LEUCADENDRON \*. Lepidocarpodendron Boerh. II.

p. 183. 184. 185. 186. 187. 188. 189. 190. 191. 194.

Hypophyllocarpendron Boerh. II. p. 205. 206. Cono-

carpodendron Boerh. II. p. 196. 199. Cinnamomea. Petal. Scalyrostrata. heom.

~~cal. quinque lacinia. l. t. 3. longa. cap. membranacea vel lacinia. l. p. 1. petalum. petalum. propria. o.~~

CAL. Perianthium commune, imbricatum : Squamis inæqualibus, persistentibus ; figura & proportione variis. propria. o.

COR. Universalis uniformis convexa.

Propria dipetala, oblonga, externe pubescens.

Petalum superius Ungue longo linearis. Limbo lanceolato indi-

viso inferne arcte adnexo petalo inferiore.

Petal. inferius Ungue longo, linearis, triplo latior.

Limbo oblongo, semicylindraceo, trifido.

STAM. Filamenta quatuor, subulata, brevissima, inserta limbo corollæ intra singulam laciniam singula.

Anthera erecta, tubulata, ex quatuor propriis in cylindrum connatis constans, oblonga.

PIST. Germen oblongiusculum desinens in Stylum subulatum longissimum. Stigma simplex.

PER. nullum. Calyx vix mutatus.

SEM. solitaria, subrotunda, coronata pilis.

REC. nudum vel villosum.

OBS. datur species calyce unifloro, non communi,

~~flor laterali extenuata ex id. ac. portion. nonum liliacum nigr. gracil.~~

86. PROTEA †. Conocarpodendron Boerh. II. p. 195. ~~Conocarpodendron~~.  
197. 198. 200. 202. 203. 204. ~~Conocarpodendron~~.

CAL. Perianthium commune, subrotundum, imbricatum : squamis subovatis. propria. o.

COR. Universalis uniformis, calyce paulo longior.

Propria monopetala : Tubo longitudine calycis ; Limbo quadrifido, glabro, patente, longitudine tubi, æquali.

STAM. Filamenta quatuor, setacea, corolla longiora. Antheræ incumbentes.

PIST. Germen subrotundum. Stylus setaceus, erectus, longitudine staminum. Stigma obtusum.

PER. nullum. Calyx immutatus.

SEM. solitaria, subrotunda, nuda.

REC. conicum, paleis distinctum.

~~flotta oblonga imbr.  
flor vestitilla. elegat.~~

*Kk. 87. CEPHALANTHUS* †. *Platanocephalus* *Vaill.* A. G.

1722. vide HM. III: 33. foliante cipit. plane  
na. Specie selenij Khasi Derta addit. uncapitata in flore et staminis. bonalid opp.  
cyp. foliop. cinnatum petalo luteo. lvs. glab. lvs. glab. lvs. glab. lvs. glab. lvs. glab.

**CAL.** Perianthium commune Receptaculo communi flores plures in capitulum colligens. Nullum in Ross. incommunis. Aperta caerulea stipe.

*Perianthium proprium*, <sup>multum in Hoff.</sup> quadripartitum, erectum, persistens.

**COR.** Petalum unicum. Tubus angustus. Limbus quadripartitus,  
longitudine tubi, reflexus, acutus. face rotunda at spiculae

**S T A M.** Filamenta quatuor, corollæ inserta, limbo<sup>n2</sup> breviora. An-  
theræ simplices. *arat. apode in dentitione aberto.*

Pist. Germen infra florem. Stylus simplex. Stigma simplex.

~~P. R. Capsula oblonga, unilocularis. (plures sinal constituunt~~

~~Spec. 1. medie P. E. R. Capsula oblonga, unilocularis. (plures sphaeræ constituant  
lunulæ ex pustulis capitulum globosum).~~ Capula <sup>2 lunc</sup> rotunda in Hoff. <sup>4 valv.</sup> Cirrula septata <sup>5 val.</sup>  
~~Spec. 2. plurima, oblonga.~~ Receptac. in Hoff. paleis 3 menses aperte latioribus  
ovatis insper. auctius pectus <sup>undulatum</sup> diversitatem.  
O B S. unitas saepe additur partibus floris. 10 cuncta celia sing. ovarium.  
fl. opp. et 3 m.

88. GLOBULARIA \*. GLOBULAIRE. *Tournef.* 265.

*Alypum Niss.* *A. G.* 1712, *t.* 18. *Dill. gen.*

**CAL.** Perianthium commune imbricatum, longitudine flosculorum:

*Perianth. proprium monophyll*

**COR. universalis fere æqualis.**  
Propria monopetala, basi tubulata, Limbo quadripartito : Laci-  
nia superiore minima reflexa, reliquis tribus inferioribus, majori-  
bus, æqualibus, labium inferius constituentibus.  
**S T A M.** Filamenta quatuor, simplicia, longitudine corollulae. An-  
theræ distinctæ.

P I S T. Germen ovatum. *Stylus simplex*, longitudine staminum.  
*Stigma obtusum* <sup>subreniforme</sup> 1. rubrum levigatum 2. <sup>luteum</sup> luteum.

P E R. nullum. *Calyx proprius connivens*, includens semen. *Receptaculum commune oblongum*, paleis distinctum.

S E M. solitaria, ovata.

~~Dipsacus~~ 89. DIPSACUS \*. CHARDON A FOULON. Tournef. 265.

Vaill. A. G. 1722. *Mala* 200

**C A L.** *Perianthium* ~~commune~~, *multiflorum*, *polyphyllum*: *foliolis* tenuibus, *flosculis* longioribus, *laxis*, *persistentiibus*.

*Perianthium proprium* vix manifestum, germini insidens. *Cor. singula monopetala. Petalum totum tubulatum rectum: ora*

*quadrifido, erecto: lacinia exteriori majori, acutiori.*

**S T A M.** *Filamenta* quatuor, capillaria, corollâ longiora. *Antheræ* incumbentes.

P I S T. Germen infra receptaculum proprium flosculi. Stylus filiformis, longitudine corollæ. Stigma simplex <sup>et lat.</sup>

P E R. nullum. RECEPTACULUM *commune* *conicum*, paleis longioribus *distinctum*.

*S E M. solitaria, columnaria, calyculo marginaceo integro coronata.*

*Calligrapha h. Purpurea* hindsh.  
a bor. fl. opp. round tenu.  
flor. radicell. ad late  
calyx 1.7th. 4 for. - inflexus ag  
cor. 1 pet. 2 ad 5. ap.  
st. 4. 4 ad 5. bractab. cor. inv. ag.  
ov. ad 1. styl. 1. ova 2.  
folia 4 pinnat. 6. sp.

cooperation. objects general go. SCA-  
S. general command Kreidek. 191  
Cal. 1 ph. 4. for info. 4. states  
encl. part 4. states states  
of 4.  
over 1.  
Capt. J. L. Lovett - polity  
other name from. (see also) (see also)

fol. opp. plorate. altero attacq. lateri insidente sic ⑩

## TETRANDRIA MONOGYNIA.

*radice apice tumida et granularis*

33

90. SCABIOSA \*. SCABIÈUSE. Tournef. 263. 264. Vaill:

A. G. 1722. Astrocephalus Vaill. ibid. Succisa. MORS DU DIABLE. Vaill. ibid. Pterocephalus Vaill. ibid.

*modus. florum nullum nisi quama sub sing. flore 1. quorum exteriores involucrando longiori ordinis. ovarium protracta signata & involucrum pro posteriori calix partite 1 phyllo supra hinc membranaceae inclusum.*

CAL. Perianthium commune multiflorum patens, polyphyllum; foliolis seriebus variis receptaculum cingentibus eique insidentibus, quorum interiora gradatim minora.

Perianthium proprium ~~duplex~~, utrumque germini insidens, quo-

rum

P. exterior brevius, membranaceum, plicatum, persistens. *ovis non intus*

P. interior quinquepartitum: *lacinis* subulato-capillaceis. *ad ipsum ampliata sed non ultra agni*

COR. singula monopetala. Petalum erectum, tubulatum, sensim ampliatum, semi-quinquefidum (quadrifidumve:) laciinis exterioribus majoribus.

STAM. Filamenta quatuor, subulato-capillaria, debilia. Antheræ *att. excalycis* oblongæ, incumbentes.

PIST. Germen infra receptaculum proprium flosculi, propriâ vaginâ involutum tanquam calyculo. Stylus filiformis, longitudine corollæ. Stigma obtusum, oblique emarginatum.

PER. nullum. RECEPTACULUM commune convexum, foliolis calycinis corollas distinguenteribus.

SEM. solitaria, ovato-oblonga, involuta, calycibus propriis coronata. *calyx membranaceus ex parte tenuis, sinuatus, convexus, crassior, rotundat. apicem acutum.*

OBS. Figura in hocce genere maxime inconstans est. Corollulæ exteriiores saepe maiores, magis inæquales. Seminum coronæ variant faciem ferunt.

## 91. KNAUTIA \*. Lychni-Scabiosa. Boerh. Linnaeum. S. yeb.

*multicolor* *Weyden. s. A. C. flat. 1. lodd. 1. lodd.*

CAL. Perianthium commune flosculos in simplicem orbem digestos continens, simplex, cylindraceum, oblongum, erectum: laciiniis subulatis, approximatis, numero flosculorum.

Perianth. proprium minimum, coronans germen. *magno inter brevi ipsius*

COR. universalis æqualis: propria monopetala, inæqualis, cujus petali *Tubus* longitudine calycis. Limbus inæqualis, quadrifidus; laciina exteriore majore, ovata.

STAM. Filamenta quatuor, tubo corollæ longiora, receptaculo inserta. Antheræ oblongæ, incumbentes.

PIST. Germen infra receptaculum proprium flosculi. Stylus filiformis, longitudine staminum. Stigma *crassiusculum*, bifidum. *in floribus I*

PER. nullum. RECEPTACULUM commune vix notabile, planum, nudum.

SEM. solitaria, *tetragona*, apice *villoso*, *coria* *brevi* *integri* *sericea* *anulata* *int* *ad* *pro* *basim*.

OBS. Calyx tubulato & simplici flosculorum orbita à Scabiosis distin-

guitur, quibus valde affinis. *calyx tubulatus* *simplici* *flosculorum* *orbita* *ad* *basim* *int* *apicem* *obtuse* *gibbosus*

*fructuosa*. *fl. lutea*. *Willd. 1. lodd.* *caulis* *ultra* *rotundat.* *caulorum* *rotundat.*

*Cal. 4 phyllo. 2 mag. 2 mag. 2 mag. 2 mag.* *Cor. 4 phyllo. 4 phyllo. 4 phyllo.*

*Cor. 4 phyllo. 4 phyllo. 4 phyllo. 4 phyllo.* *ovaria 1.* *ovaria 1.* *ovaria 1.* *ovaria 1.*

*fructuosa*. *fl. lutea*. *Willd. 1. lodd.* *caulis* *ultra* *rotundat.* *caulorum* *rotundat.*

*Cal. 4 phyllo. 4 phyllo. 4 phyllo. 4 phyllo.* *Cor. 4 phyllo. 4 phyllo. 4 phyllo. 4 phyllo.*

*Cor. 4 phyllo. 4 phyllo. 4 phyllo. 4 phyllo.* *ovaria 1.* *ovaria 1.* *ovaria 1.* *ovaria 1.*

*fructuosa*. *fl. lutea*. *Willd. 1. lodd.* *caulis* *ultra* *rotundat.* *caulorum* *rotundat.*

— Portulaca L. M. & 257. Baroni. Barth. Hata hant. Haast. Haad. Belg. En hin aleamone  
arborescens ramos ligulis opp. glandul. fl. opp. alcavus concrevata sed parallelo ordin-  
flo corimb termin. anig. cal. ovar. inic. 4 p. p. cor 1 pet. tubul. 4 W. st. 4. urid.  
**TETRANDRIA MONOGYNIA.**

**TETRANDRIA MONOGYNIA.**  
Bracts 34 orat long. globul. polyapp. uncinis. ?/tip. caudat. exst. Tangaraceae ordin. <sup>simplicis,</sup>  
cavilea.

- 92. *SHERARDIA* \*. Dill. gen. 3.  
*fl. pyriforme verticillata. fl. anelli min.*  
 CAL. *Perianthium parvum, quadridentatum, germini insidens,*  
*persistens.*  
 COR. *Petalum unicum. Tubus cylindraceus, longus. Limbus qua-*  
*dripartitus, planus, acutus.*  
 STAM. *Filamenta quatuor, ad apicem tubi posita. Antheræ sim-*  
*plices.*  
 PIST. *Germen didymum, oblongum, infra receptaculum. Stylus*  
*filiformis, superne bifidus. Stigmata capitata.*  
 PER. *nullum. Fructus oblongus, coronatus, longitudinaliter in*  
*duo semina separabilis.*  
 SEM. *bina, oblonga, apice tribus acuminibus notata, hinc conve-*  
*xa, inde plana.*

93. fol. 4 ad 8. fls. 6 wh. tenui pedunculati. **ASPERULA** \*.

- C A L. *Perianthium parvum, quadridentatum, germini insidens.*  
C O R. *Petalum unicum. Tubus cylindraceus, longus. Limbus quadripartitus: laciniis oblongis, obtusis, reflexis.*  
S T A M. *Filamenta quatuor, ad apicem tubi posita. Antheræ simplices.*  
P I S T. *Germen didymum, subrotundum, infra receptaculum. Stylis filiformis, superne bifidus. Stigmata capitata.*  
P E R. *Baccæ siccæ, duæ, globosæ, coalitæ.*  
S E M. *solitaria, subrotunda, magna.*

-94. SPERMACOCE †. Dill. elth. 277.

- C A L. *Perianthium* parvum , quadridentatum , germini insidens ,  
persistens.

C O R. *Petalum* unicum. *Tubus* cylindraceus , calyce longior. *Lim-  
bus* quadripartitus , patenti-reflexus , obtusus.

S T A M. *Filamenta* quatuor , subulata , corollâ breviora. *Antheræ*  
simplices.

P I S T. *Germen* subrotundum , compressum , infra receptaculum.  
*Stylus* simplex , superne bifidus. *Stigmata* obtusa.

P E R. *Capsulæ* duæ , connatæ , oblongæ , hinc gibbæ , inde planæ ,  
obtusæ , singula bicornis.

S E M. solitaria , subrotunda.

95. CRUCIANELLA \*. Rubeola Tournef. 50.

- CAL. *Perianthium* <sup>*scapinum*</sup> *triphyllum*: *foliolis* lanceolatis, carinulatis,  
acuminatis, rigidis, conniventi-compressis.  
COR. *Petalum* unicum. *Tubus* cylindraceus, filiformis, calyce  
longior. *Limbus* quadrifidus, parvus.  
STAM. *Filamenta* quatuor, in ore tubi posita. *Antheræ* simplices.

— *Carex corylifolia* Drom. p. 144. 2. fig. 2. —  
caesp. stiff. lvs. 1-2 ft. long. lvs. 1-2 ft.  
wide. Culms con. lvs. rect. blade 4 ft.  
Copeat. st. 1 ft. culm apic. tub. linear. alt. ex lvs  
Brown. ov. 1. styl 1. fig. 2. long & cyl. green  
rect. in fig. base glob. inflate 2. lvs. polyphy. oblong  
Dense. lvs. narrow. obtuse. Disseminate after  
fol. opp. time. ? lvs. caesp. con. volv. Brown.

# TETRANDRIA MONOGYNIA.

PIST. Germen compressum, inter calycem & corollam. Stylus filiformis, longitudine tubi. Stigmata duo, obtusa, unicum stylum terminans, alterum infra apicem styli sessile.

P. E. R. Capsulæ duæ, connatæ.

*S. M. solitaria, oblonga.*

—96. HOUSTONIA †. Paronychiæ affinis Moris. III. f. 15.

t. 4. f. 1. auctore Clariss. D. Gronovio.

CAL. Perianthium minimum, quadridentatum, erectum, persis-  
stens. *inflorescens* tubo brevior.

*Cor. Petalum unicum, infundibuliforme. Tubus cylindraceus,  
longus. Limbus quadripartitus, patens, laciinis subrotundis.*

**S T A M.** *Filamenta* quatuor, in collo corollæ, minima. *Antheræ* simplices.  
**P I S T.** *Germen* subrotundum, compressum. *Stylus* simplex, staminibus brevior. *Stigma* bifidum, acutum.

P E R. Capsula subrotunda, didyma, bilocularis, bivalvis, valvulis  
dissepiamento oppositis. 3/10.

**S E M.** solitaria.

—97. GALIUM \*. CAILLELAIT. Gallium. Tournef. 39.

fl. 4 ad 10. fl. crypt

C A L. *Perianthium* minimum, quadridentatum, germini insidens.

**COR.** Petalum unicum, quadripartitum, planum, acutum, tubo nullo.

STAM. Filamenta quatuor, subulata, corollâ breviora. Antheræ simplices.

Pist. Germen didymum, infra receptaculum. Stylus filiformis,  
semi-bifidus, longitudine staminum. Stigmata globosa.

P E R. Baccæ siccæ, duæ, globosæ, coalitæ, non adhærentes.

**S E M.** *solitaria*, *reniformia*, *magna*.

—98. APARINE \*. GRATTERON. *Tournef.* 39.

CAL. Perianthium minimum, quadridentatum, germini insidens.

**COR.** *Petalum unicum, quadripartitum, planum, acutum, tuba nullo.*

STAM. Filamenta quatuor, subulata, corollâ breviora. Antheræ simplices.

P I S T. Germen didymum, infra receptaculum. Stylus filiformis,  
semi-bifidus, longitudine staminum. Stigmata capitata.

PER. Baccæ siccæ, duæ, globosæ, coalitæ, villis uncinatis, rigidis adhærentes.

S E M. solitaria, subrotunda, umbilicata, magna, <sup>ad monogrammam ac</sup>  
~~mitchillae~~ mitchillae <sup>et</sup> <sup>2</sup> <sup>4</sup> <sup>8</sup> <sup>12</sup> <sup>16</sup> <sup>20</sup> <sup>24</sup> <sup>28</sup> <sup>32</sup> <sup>36</sup> <sup>40</sup> <sup>44</sup> <sup>48</sup> <sup>52</sup> <sup>56</sup> <sup>60</sup> <sup>64</sup> <sup>68</sup> <sup>72</sup> <sup>76</sup> <sup>80</sup> <sup>84</sup> <sup>88</sup> <sup>92</sup> <sup>96</sup> <sup>100</sup> <sup>104</sup> <sup>108</sup> <sup>112</sup> <sup>116</sup> <sup>120</sup> <sup>124</sup> <sup>128</sup> <sup>132</sup> <sup>136</sup> <sup>140</sup> <sup>144</sup> <sup>148</sup> <sup>152</sup> <sup>156</sup> <sup>160</sup> <sup>164</sup> <sup>168</sup> <sup>172</sup> <sup>176</sup> <sup>180</sup> <sup>184</sup> <sup>188</sup> <sup>192</sup> <sup>196</sup> <sup>200</sup> <sup>204</sup> <sup>208</sup> <sup>212</sup> <sup>216</sup> <sup>220</sup> <sup>224</sup> <sup>228</sup> <sup>232</sup> <sup>236</sup> <sup>240</sup> <sup>244</sup> <sup>248</sup> <sup>252</sup> <sup>256</sup> <sup>260</sup> <sup>264</sup> <sup>268</sup> <sup>272</sup> <sup>276</sup> <sup>280</sup> <sup>284</sup> <sup>288</sup> <sup>292</sup> <sup>296</sup> <sup>300</sup> <sup>304</sup> <sup>308</sup> <sup>312</sup> <sup>316</sup> <sup>320</sup> <sup>324</sup> <sup>328</sup> <sup>332</sup> <sup>336</sup> <sup>340</sup> <sup>344</sup> <sup>348</sup> <sup>352</sup> <sup>356</sup> <sup>360</sup> <sup>364</sup> <sup>368</sup> <sup>372</sup> <sup>376</sup> <sup>380</sup> <sup>384</sup> <sup>388</sup> <sup>392</sup> <sup>396</sup> <sup>400</sup> <sup>404</sup> <sup>408</sup> <sup>412</sup> <sup>416</sup> <sup>420</sup> <sup>424</sup> <sup>428</sup> <sup>432</sup> <sup>436</sup> <sup>440</sup> <sup>444</sup> <sup>448</sup> <sup>452</sup> <sup>456</sup> <sup>460</sup> <sup>464</sup> <sup>468</sup> <sup>472</sup> <sup>476</sup> <sup>480</sup> <sup>484</sup> <sup>488</sup> <sup>492</sup> <sup>496</sup> <sup>500</sup> <sup>504</sup> <sup>508</sup> <sup>512</sup> <sup>516</sup> <sup>520</sup> <sup>524</sup> <sup>528</sup> <sup>532</sup> <sup>536</sup> <sup>540</sup> <sup>544</sup> <sup>548</sup> <sup>552</sup> <sup>556</sup> <sup>560</sup> <sup>564</sup> <sup>568</sup> <sup>572</sup> <sup>576</sup> <sup>580</sup> <sup>584</sup> <sup>588</sup> <sup>592</sup> <sup>596</sup> <sup>600</sup> <sup>604</sup> <sup>608</sup> <sup>612</sup> <sup>616</sup> <sup>620</sup> <sup>624</sup> <sup>628</sup> <sup>632</sup> <sup>636</sup> <sup>640</sup> <sup>644</sup> <sup>648</sup> <sup>652</sup> <sup>656</sup> <sup>660</sup> <sup>664</sup> <sup>668</sup> <sup>672</sup> <sup>676</sup> <sup>680</sup> <sup>684</sup> <sup>688</sup> <sup>692</sup> <sup>696</sup> <sup>700</sup> <sup>704</sup> <sup>708</sup> <sup>712</sup> <sup>716</sup> <sup>720</sup> <sup>724</sup> <sup>728</sup> <sup>732</sup> <sup>736</sup> <sup>740</sup> <sup>744</sup> <sup>748</sup> <sup>752</sup> <sup>756</sup> <sup>760</sup> <sup>764</sup> <sup>768</sup> <sup>772</sup> <sup>776</sup> <sup>780</sup> <sup>784</sup> <sup>788</sup> <sup>792</sup> <sup>796</sup> <sup>800</sup> <sup>804</sup> <sup>808</sup> <sup>812</sup> <sup>816</sup> <sup>820</sup> <sup>824</sup> <sup>828</sup> <sup>832</sup> <sup>836</sup> <sup>840</sup> <sup>844</sup> <sup>848</sup> <sup>852</sup> <sup>856</sup> <sup>860</sup> <sup>864</sup> <sup>868</sup> <sup>872</sup> <sup>876</sup> <sup>880</sup> <sup>884</sup> <sup>888</sup> <sup>892</sup> <sup>896</sup> <sup>900</sup> <sup>904</sup> <sup>908</sup> <sup>912</sup> <sup>916</sup> <sup>920</sup> <sup>924</sup> <sup>928</sup> <sup>932</sup> <sup>936</sup> <sup>940</sup> <sup>944</sup> <sup>948</sup> <sup>952</sup> <sup>956</sup> <sup>960</sup> <sup>964</sup> <sup>968</sup> <sup>972</sup> <sup>976</sup> <sup>980</sup> <sup>984</sup> <sup>988</sup> <sup>992</sup> <sup>996</sup> <sup>1000</sup>

~~Pavetta~~ *Pavetta* Brotvar p. 142. 2. 6. f 1. ~~Cult~~ 12  
Pavetta hirsuta Cal. typus Sympy. 1. phyll. turbid. parviflora hydrol.  
52. 10 Cor. 1. pet. yr. 4 folia acuta pallidiora. Ixora 2 a lata  
unisexualis et 4. stylis diversis tubi laevi alb.  
stipulae ovato-lanceolatum. style 1. obt. 2. long. connex  
calyx non glab. 1. loc. 1 sp.  
videtur glab. foli 4 lobata.  
lignosa. fol. opp. c. o. Stipulas setacea petiolata integrata

— Zandveld. h. m. 9. 2.76. @ 1. pied. in arenosa ped. stolonifera, raro d'alte 4 gradi fol. opposte  
basi ciliatae rufa (caulis). caulinis ciliatae granulosa ex. flm. 2 ad 3 in long. ala opp. scabia.  
numeris nullis tato long. 6 cm. gatid 4.63. sq. 0.64. med. 0.67. 2 pd. cap. 2 lobi 2 granulatis long.  
verbena - genus diffusa s. gl. 67 **DRIA MONOGYNIA**

# DRIA MONOGYNIA.

sum, inter calycem & corollam. *Stylus* fili-  
tubi. *Stigmata* duo, obtusa, *unicum stylum*  
*infra apicem styli sessile.*

Cat. mon. 2 fidei ag.  
Con. mon. 5. ineq. rinv  
Sta. A. authoris ineq. verus apud. tali ad dico  
alit. dubit.  
Stigma 1. violaceum. puncto erasum  
Rugos. gress. orbicularis planus.  
Serr. 2. conica recta inad. lassimis  
tubo per.

*Stereodia*. Sgl. 171. 1. *verbena* spicata angustifolia  
2. *anthidif.* 2. *Stereodia*  
semi-oblonga  
spica terminalis oblonga  
folia velutinacinate.

†. Paronychiæ affinis *Moris*. III. f. 15.  
auctore Clariss. D. Gronovio.

*Enchenopis cognitata* calice supero. El alto  
gasteris.

CAL. Perianthium minimum, quadridentatum, erectum, persis-  
stens. *infra florum tubo bivalve*

**COR.** *Petalum unicum*, infundibuliforme. *Tubus cylindraceus*,  
*longus. Limbus quadripartitus*, patens, *laciniis subrotundis*.

STAM. Filamenta quatuor, in collo corollæ, minima. Antheræ simplices.

Pist. Germen subrotundum, compressum. Stylus simplex, staminibus brevior. Stigma bifidum, acutum.

P E R. Capsula subrotunda, didyma, bilocularis, bivalvis, valvulis  
dissepiamento oppositis. 21p.

**S E M. solitaria.**

—97. **GALIUM** \*. CAILLELAIT. Gallium. *Tournef.* 39.

fl. 4 ad 10. fl. cypria

C A L. *Perianthium minimum*, *quadridentatum*, *germini infidens*.

**COR.** *Petalum unicum*, quadripartitum, planum, acutum, tubo nullo.

STAM. *Filamenta* quatuor, subulata, corollâ breviora. *Antheræ* simplices.

P I S T. Germen didymum, infra receptaculum. Stylus filiformis,  
semi-bifidus, longitudine staminum. Stigmata globosa.

P E R . Baccæ siccæ , duæ , globosæ , coalitæ , non adhærentes .  
S E M . solitaria , reniformia , magna .

—98. APARINE \*. GRATTERON. *Tournef.* 39.

C A L. *Perianthium* minimum, quadridentatum, germini insidens.

**COR.** Petalum unicum, quadripartitum, planum, acutum, tubo nullo.

STAM. Filamenta quatuor, subulata, corollâ breviora. Antheræ simplices.

P i s t. Germen didymum, infra receptaculum. Stylus filiformis semi-bifidus, longitudine staminum. Stigmata capitata.

PER. Baccæ siccæ, duæ, globosæ, coalitæ, villis uncinatis, rigidis adhærentes.

*S E M. solitaria, subrotunda, umbilicata, magna ad monitum ac  
mitula sive 4 deponit.*

*Pavetta Brownii* p. 142. 2. S. f. 1.  
Pavetta brownii Cal. typ. super. 2. phyll. rubra. perianth  
52. 10 Cor. 1. pub. 7r h. l. acute. jasminii Ix.  
unisexualis st. h. l. liliifl. corolla tub. laev. att.  
stipitata ovary reflexum. styl. 1. obt. 2. long. connex  
calyx 5lob. 1. loc. 1/4.  
widet 5em 1. glob. foli. 4 lobus.  
stipitata Sol. opp. cymosa. stipulae setacea petio





fol. alt. glapp. utryg. lat. involuta. Et adhuc castig. datus.

三八

TETRANDRIA MONOGYNYA.  
*et unigata et pauciflora.*

*Plantago monanthos*, Radin aperte unisata et praemorsa.  
aproximatae sordi lutea.

~~monica at~~ 106. PLANTAGO\*. PLANTAIN. Tournef. 48. Coronopus  
~~flavus hercynicus~~ Tournef. Psyllium. HERBE AUX PUCES. Tournef. Gramen  
~~et majoralis~~ junc; 30. Moris. III. s. 8. t. 9. f. ult.  
~~andreae pedunculata~~  
~~carr. h. phyt.~~

**PL**a in plant. latif. inanis Cal. exserta 3 phyll. vag. fol. 2. lateribus non rotundatis. crenulatis et 1. Dorsali plana, crenulata.  
ventrali nulla sed aperte fuso forte 4<sup>o</sup> quama dorsali appressa clavis dorsali, velut duriorate q. calyx.  
**in Plantag. latif.** **CAL.** Perignthium quadripartitum, erectum, brevissimum, perfi-  
stens. <sup>hac 4<sup>o</sup> quama loco flori iugis in spicam imbutatum collecta.</sup>  
**COR.** Petalum tabescens. Tubus cylindraceo-globosus. Limbus  
quadripartitus, depresso : laciniis ovatis, acutis.  
**STA M.** Filamenta quatuor, capillaria, erecta, omnium longissima.

*P. T. A. M. Filamenta quatuor, capillaria, erecta, omnium longissima.  
Antheræ oblongiæ, compressæ, incumbentes. Plantæ latifloraæ.  
P. I. S. T. Germen ovatum. Stylus filiformis, staminibus dimidio bre-  
vior. Stigma simplex.*

PER. Capsula ovata, bilocularis, horizontaliter dehiscens, dissepi-  
mento libero. ~~Fructus~~ capsula <sup>centrale</sup> leviter <sup>modice</sup> crassius. velutina latula  
<sup>et</sup> tenuis. <sup>et</sup> levigata. velutina. <sup>et</sup> levigata.  
SEM. plurima, oblonga. <sup>versus</sup> <sup>versus</sup> <sup>versus</sup> <sup>versus</sup> <sup>versus</sup> <sup>versus</sup>  
OBS. Calyx in quibusdam inaequalis est, in aliis aequalis.

107. SANGUISORBA <sup>*Sangu. col.*</sup> \*. Rupp. Pimpinellæ species

*Pimpinella gal. da. 868.*, capite dorsi PIMPRENELLE. Tournes. famili. Rosaceen fl. spirali densis long. quamq. 2. lippett. postea que sive rotat calicem caputum sellata.

**CAL.** Perianthium diphyllum: foliolis oppositis, brevissimis, caducis.  
**COR.** Petalum planum, quadripartitum; lacinias ovatis, obtusis, unguibus cohaerentibus. Sali gne late et rotundata, parvula.  
**STAM.** Filamenta quatuor, sc

**S I A M.** *Filamenta* quatuor, superne latiora, longitudine corollæ.  
*Antheræ* subrotundæ, parvæ.  
**P I S T.** *Germen* quadragonum, inter calycem & corollam positum.  
*Stylus* filiformis, brevissimus. *Stigma* obtusum.  
**P E R.** *Capsula*, parva, bilocularis.  
**S E M.** parva. Cap. 1. long. 15 p. in grec. canadum. tunc calyx. int. ova. testa  
est. testa stylis ac semina. Embryo rectus. glob. 2. c. styl. endosperm. alter crassus. & apice in creu-  
s. per ovarium. radicell. R. radicell. calyx. petenti. O. O. C. D.

— 108. ~~CENTUNCULUS~~<sup>\*.</sup> Dill. gen. s. *Anagallidastrum*

~~188. CENTUNCULUS~~ +. Dill. gen. s. *Anagallidastrum*  
~~1. panic flw. in whorls~~ Mich. 18. *Anagallidis* Sp. *Vaill.* B. P. 4. f. 2.  
~~alt. cymes~~ ~~sp. 1-2 bracts~~ ~~whorl~~ ~~oblique~~ ~~reddish~~ ~~green~~ ~~yellow~~ ~~alt.~~

**CAL.** *Perianthium quadripartitum*, patens, persistens: *laciiniis acutis*, lanceolatis, corolla longioribus.  
**COR.** *Petalum unius*.

**COR.** *Petalum unicum*. *Tubus* <sup>un</sup>*globo**sus*. *Limbus* *quadripartitus*, *planus*: *lacinias* *subovatis*.

STAM. Filamenta quatuor, longitudine fere corollæ. Antheræ simplices.

Pist. Germen subrotundum, intra tubum sessile. Sili. ang.

*Capsula globosa*, unilocularis, horizontaliter dehiscens.  
*SEM.* plurima, subrotunda, minima.

DE M. plurima, subrotunda, minima. *Aegyptiantele angustata*

*Malvaceae*. Lin. + Crotalariaeum. Brovnik 145. 2. 1. *Croton* <sup>l</sup> *luteus*: 4 ad 5 per

— 1114. *Rhabdonia*. Lin. & *Crocosmia* — *Pavonia* 1115. L. 17. filypoda Buderus. 4 ad 3 per  
cat. flower. Lin. cat. sp. ph. 4 per ad. ph. per. infery. Con. sp. ph. per. bracts.  
8. *Centaurium* et. L. Digit. calyx tubo concreta in 2. 00 ad 1. styl. brevis. dig. 8. obt.  
major 2g. hirsutus caps. glob. 16m. 1.5mm. Linn 1. glob.  
fol. ovv. veronicae seu aplolethera.

foli opp. veronicae seu apocynaceae  
caule rotundato opp. ciliata  
flav. ~~longit. corollis~~ allatilis, ligustrinum, long. fl. virides

fol. 20d. pl. pl. 20. 9. 10. involuta sic an hier 6  
Ex me dimidiatim pl. pl. et circinaliter, cum pedicello involuta et recepta intra cuneum  
quid foliis late petiolis connivit format. Stip. o.  
**TETRANDRIA MONOGYNIA.** fol. punctata hyperdi  
fl. / pedic. non nullus sed an medio petioli conniv. flor ut in chrysophyllum ang. et canto, quam 1 stip.  
39  
**EPIMEDIUM \***. Tournef. 117. oddius et fumarial  
109. Siz. D. 1. fl. et fructu cari. gnetos ad oliveras subtilis.  
et pascuas

CAL. Perianthium tetrphyllum : foliolis ovatis, obtusis, concavis, patentibus, parvis, petalis directe (non alternatim) subjetis, caducis.

*Cor. Petala quatuor, ovata, obtusa, concava, patentia.*  
*Nectaria quatuor, <sup>racemata</sup> cyathiformia, fundo obtuso, magnitudine pe-*

*Nectaria quatuor*, cyathiformia, rando obtuso, magnitudine petalorum, petalis <sup>elongata</sup> incumbentia, margine oris affixa receptaculo.  
STAM. Filamenta quatuor, subulata, stylum prementia. Antheræ elongatae, erectæ, biloculares, bivalves, à basi apicem versus de-

oblongæ, erectæ, biloculares, bivalves, a bâli apicem verius de-  
hiscentes, dissepimento libero. *ut Barbæ cui adest*  
Pist. Germen oblongum. Stylus germine brevior, longitudine  
staminum. Stigma simplex.

staminum. Stigma simplex.  
P. E. R. Siliqua oblonga, acuminata, unilocularis, bivalvis.

P E R . Siliqua oblonga , acuminata , unilocularis , bivalvis .  
S E M . plurima , oblonga .

SEM. *plumaria*, oblonga.

110. CORNUS \* CORNOUILLER. Tournef. 410. Virga xavia R. B.  
Sum. 2. annis interiore⁹ rameal antiqua gerunt. pili naurulare⁹.  
Sanguinea SANGUEN. Dill. Cornus habacea. Mesomora R.  
Cornulus heist. cornus foec. - sanguinea involucro nullo. pili naurulare⁹  
CAL. Involucrum tetraphyllum, commune, multiflorum, pedun-  
culi-ferum; foliolis ovatis, oppositis minoribus, coloratis, deci-  
duis. flors umbellate par compact. ramis tricus. ante fl. flores. involucrum nullum in cornufloro.  
Perianthium minimum, quadridentatum, germini insidens, de-  
ciduum.

*Ciduum.*  
COR. Petala quatuor, oblonga, acuta, plana, involucro minora. *dorsata. flos*  
*innuptus. alt-*  
*erecta. corollâ longiora.* *alii in lan-*

**COR.** Petala quatuor, oblonga, acutæ, p.  
**STAM.** Filamenta quatuor, subulata, erecta, corollâ longiora.  
Antheræ subrotundæ, incumbentes. . Quandam primitiæ per se latitudine inveniatur in rarae solent latitudine. sibi  
succincta subulatum Stylus filifor-

P I S T. Germen subrotundum, infra receptaculum. Stylus filiformis, longitudine corollæ. Stigma obtusum.

mis, longitudine coronæ. Stigma obtusum.  
P. Drupa subrotunda, umbilicata. Nucleus oblongus.

**S E M.** *Nux cordata* seu *oblonga*, *bilocularis*. *Nucleus oblongus.* *ad 6*  
*Hanc malum: foliacea lateribus leviter involuta, sic conniventer.*

**EUONYMUS** \*. FUZAIN. Tournef. 388.

**III.** EUONYMUS. FUZAIN. *varius*. *folia* *opposita* *quadrangula* *lacinia* *parallelia* *folia* *opposita* *lacinia* *parallelia* *folia* *opposita* *lacinia* *parallelia*

**CAL.** Perianthium monophyllum, quadripartitum, planum: laci-  
folia in parallela

*Petala quatuor, ovata, plana, patentia, calyce longiora.*

**STAM.** Filamenta <sup>versatiles</sup> quatuor, subulata, erecta, corollâ breviora, calvi opp.  
Antheræ didymæ. <sup>versatiles</sup>

*Stam.* Filamenta quatuor, rufata, germini tanquam receptaculo imposita. Antheræ didymæ. *Anthr.* *Stylus* *luminatum*. *Stylus brevis*, simplex. *Stigma obtusum*.

Pist. Germen acuminatum. Stylus brevis, simplex. Stigma obtusum.

*sum.*  
P. *Capsula succulenta, colorata, quadragona, quadrangularis,*

PER. Capsula succulenta, colorata, quadrangulata,  
quadricuspis, quadrilocularis, quadrivalvis.  
Semina granulata obvoluta.

*S E M. solitaria, ovata, calyprâ obvoluta.*

**O**bs. Fructificatio, in quibusdam inavians vel speciebus annis  
quintam numeri partem.

quintam numeri partem.  
Quibusdam Filamenta, præterquam acumina receptaculi, nulla.

St. li. longe. capi. glob. 1 lb. 2 valer 1/16. Count 1. glob. uero.  
Eli. sicut aputat' a tunc condato suuitate ceteri.

