



Hunt Institute for Botanical Documentation
5th Floor, Hunt Library
Carnegie Mellon University
4909 Frew Street
Pittsburgh, PA 15213-3890
Telephone: 412-268-2434
Email: huntinst@andrew.cmu.edu
Web site: www.huntbotanical.org

The Hunt Institute is committed to making its collections accessible for research. We are pleased to offer this digitized item.

Usage guidelines

We have provided this low-resolution, digitized version for research purposes. To inquire about publishing any images from this item, please contact the Institute.

About the Institute

The Hunt Institute for Botanical Documentation, a research division of Carnegie Mellon University, specializes in the history of botany and all aspects of plant science and serves the international scientific community through research and documentation. To this end, the Institute acquires and maintains authoritative collections of books, plant images, manuscripts, portraits and data files, and provides publications and other modes of information service. The Institute meets the reference needs of botanists, biologists, historians, conservationists, librarians, bibliographers and the public at large, especially those concerned with any aspect of the North American flora.

Hunt Institute was dedicated in 1961 as the Rachel McMasters Miller Hunt Botanical Library, an international center for bibliographical research and service in the interests of botany and horticulture, as well as a center for the study of all aspects of the history of the plant sciences. By 1971 the Library's activities had so diversified that the name was changed to Hunt Institute for Botanical Documentation. Growth in collections and research projects led to the establishment of four programmatic departments: Archives, Art, Bibliography and the Library.

On Elaioplasts.

BY

RUDOLF BEER, B.Sc., F.L.S.

With Plate IV.

IN 1888 Wakker (1) described the occurrence of a conspicuous body lying in the epidermal cells of the leaves and in the superficial tissues of the root and stem of *Vanilla planifolia*. This body is somewhat larger than the nucleus and considerably larger than the amyloplasts; it possesses a sharply-defined outline and a peculiar, somewhat yellowish colour. In each plate-like epidermal cell one such body occurs, and it often lies near the nucleus, although in other cases it may occupy a different position in the cell. As this body consists of protoplasmic material and contains oil, Wakker named it *Elaioplast*, or oil-former.

He made a careful study of the effect of reagents upon the elaioplast, and briefly described the gradual disappearance of these bodies in older cells. Wakker was unable to study the origin of the elaioplasts as he had no suitable material for this purpose. He found elaioplasts also to occur in the cells of another species of *Vanilla*, known to him under the name of *Vanilla aromatica latifolia*. In 1893 Zimmermann (2) found similar bodies to occur in *Funkia coerulua*, *F. lancifolia*, *F. Sieboldiana*, *Dracaena* sp., *Ornithogalum scilloides*, *Agave americana*, *A. Mitis*, and in *Oncidium suave*.

Raciborski (3) in the same year described elaioplasts in the tissues of various species of *Ornithogalum*, *Albuca*, *Funkia*, and *Gagea*.

Zimmermann further found these bodies in the internal cells of the stem of *Psilotum*, and in the perianth leaves of *Maxillaria picta*. The shape of the elaioplasts differs in various plants, but it is usually constant in the same species. Spherical forms, grape-like bodies, irregular plasmodium-like masses, have all been described.

Usually only one elaioplast occurs in a cell, but in some cases they may be more numerous (e. g. *Ornithogalum*). The finer structure of the elaioplast has been carefully examined both by Zimmermann and by Raciborski. It has a finely granular appearance due to the occurrence of a number of tiny, highly refractive spherical bodies lying in its substance. One or more less refractive spots often occur within the elaioplast. The elaioplasts of

the species of *Gagea* are said by Raciborski to possess a somewhat different structure. In these plants the spherical elaioplasts are characterized by the feeble development of the stroma: they are surrounded by a plasmatic envelope and contain within this an oily substance like that found in other elaioplasts. The micro-chemical observations of Wakker and Zimmermann have shown that the elaioplasts consist of a plasmatic matrix or stroma in which are embedded numerous minute oil drops. Zimmermann points out that these oil drops show a close similarity in their reactions with the oil drops obtained from plastids.

The origin of the elaioplasts has been very little studied. Raciborski states that in *Ornithogalum umbellatum* they arise as small, highly refractive spherules which always lie at one pole of the usually elongated cell-nucleus.

Although Wakker had no material with which to work out the development of the elaioplast, he ventures an interesting suggestion at the conclusion of his account of the oil-bodies of liverworts. 'Leider ist durch diese Mittheilung der Ursprung der Elaioplasten nicht ausgemacht, es ist mir aber äusserst wahrscheinlich, dass es bei den Lebermoosen metamorphosirte Chlorophyllkörner sind. Vielleicht ist dieses auch bei *Vanilla* der Fall.'

Garjeanne's work (5) on the oil-bodies of the Jungermanniales clearly indicates that these bodies have a different origin to that suggested by Wakker; my own observations on elaioplasts recorded below will, however, show that his guess was nearer the truth in the case of some Phanerogamic elaioplasts.

The function of the elaioplast is quite unknown. Wakker believed that they might be oil-formers much as leucoplasts are starch-formers. Zimmermann offered the suggestion that they might prove to be parasitic or symbiotic fungi living within the cells of the higher plant.

Raciborski, however, considers them to be normal organs of the cell in which they occur, and classes them with oil-bodies, tannin-vesicles, and ordinary vacuoles.

In the liverworts oil-bodies have been known to occur since the time of Gottsche, and even earlier. The first really fundamental description of these bodies is due to Pfeffer (4), and quite recently their development has been fully worked out by Garjeanne (5). Oil or fat bodies of a somewhat similar appearance to those of the liverworts have been described by Radlkofer (6 and 7), Monteverde (8), Solereder (9), and others, in the tissues of a number of Dicotyledons and Monocotyledons. Opinion varies very much as to the relation of these oil-bodies of liverworts and Phanerogams to the elaioplasts. Some believe the two structures to be closely allied, whilst others are of opinion that they are radically different from one another.

From this brief survey of the literature it will be seen that elaioplasts,

corresponding to those discovered by Wakker in *Vanilla*, have up to the present been found only in Monocotyledons, with the single exception of *Psilotum* reported by Zimmermann.

Moreover, apart from a few quite insufficient observations we are still entirely in the dark regarding their true nature, their significance, their origin, and their fate.

During some observations upon the pollen-grains of Compositae I was interested to find bodies which resembled the Monocotyledonous elaioplasts occurring in the tissues of the floral region of *Gaillardia Lorenziana*¹.

This would extend the distribution of elaioplasts to a member of the Dicotyledons.

My first endeavour was to make sure that I was dealing with true elaioplasts. In the hairs upon the corolla of young flowers, where I first observed these bodies, they occur as more or less spherical, highly refractive, granular structures usually somewhat larger than the nucleus, and in most cases lying singly in a cell, although two or even more such bodies were occasionally met with.

Their reactions were found to be as follows:—

1. Heated gently upon the slide (whilst lying in a physiological salt solution), drops of oil are exuded from their surface (Fig. 13).

2. Osmic acid (as this occurs in Flemming's stronger solution) turns them black or brown.

3. Potassium bichromate after twenty-four hours' action leaves them quite colourless, but causes the extrusion of oil globules as in 1.

4. Alkannin solution in 60 per cent. alcohol colours the bodies deeply red.

5. Iodine (in KI) colours them brown and causes the extrusion of oil-globules (Fig. 12).

6. Absolute alcohol dissolves out the oil from their interior and leaves them vacuolated.

7. In strong HNO_3 (warmed), followed by NH_3 , they give the Xanthoproteic reaction (viz., deep yellow coloration).

8. Glacial acetic acid after twenty-four hours' action causes great extrusion of oil drops but no solution.

9. 10 per cent. KOH. after twenty-four hours' action dissolves neither stroma nor oil drops.

These reactions, combined with their general appearance, show that the bodies occurring in the hairs of *Gaillardia* are in all respects similar to the elaioplasts described by previous authors.

In very young hairs from capitula which were still quite small and

¹ *Gaillardia Lorenziana* is a German variety of *G. picta*, which itself appears to be only a garden variety of *G. pulchella*. I have not yet had an opportunity of examining any other form or species of *Gaillardia* except the one mentioned above.

immature and entirely enclosed within the involuclral bracts, no elaioplasts were yet to be seen. The cell contained a nucleus and cytoplasm which partly formed a peripheral layer and partly extended in strands and bars through the cell cavity (Figs. 1 and 2). Embedded in the cytoplasm was a number of small, highly refractive grains which had all the appearances of ordinary leucoplasts. That these refractive grains are really leucoplasts is confirmed by two facts.

Firstly, the resemblance between the unquestionable, starch-forming leucoplasts occurring, for example, in the hairs which cover the very young leaves, and the highly refractive grains contained in the corolla-hairs, is complete, although starch is not found in the latter under the usual conditions of growth.

Secondly, if the enveloping bracts be removed from a young inflorescence without detaching it from the parent plant, and the corolla-hairs exposed to a strong insolation, starch can be seen to have developed in some of these refractive grains.

For these reasons I believe the highly refractive grains occurring in the cells of the corolla-hairs to be leucoplasts, some of which, however, may have lost the power of starch-formation.

In somewhat older hairs these plastids, a number of which show signs of undergoing degeneration, tend to aggregate together at one or more spots within the cell. Not infrequently this aggregation of the plastids is in the neighbourhood of the nucleus, but in many instances it is found to occur at other regions of the cell (Figs. 3 and 4).

At first the aggregation of the refractive grains is a very loose one, but it gradually grows closer and closer (Fig. 5) until the compact, highly refractive bodies are formed, which we have already recognized as elaioplasts (Figs. 6, 7, and 8). The elaioplasts in the corolla-hairs of *Gaillardia* are, therefore, formed by the aggregation of plastids and their degeneration products at one or more spots in the cell. Within the elaioplast the plastids soon appear to undergo further degeneration with the production of an oily material. That Zimmermann should find a close similarity between the oil of the elaioplasts studied by him and the oil obtained from plastids is no longer surprising.

All the plastids of the cell have not clumped together within the elaioplast. A certain proportion still remain scattered through the cell (Fig. 7).

For some time there is little alteration within the cell. The conspicuous elaioplast may lie in almost any part of the cell, but often it takes up a position near the nucleus. In some instances it entirely envelops the nucleus, as I have represented in Fig. 8.

In much older hairs we find the elaioplast undergoing a change. Its outline becomes less regular, and in some cases it becomes drawn out

and elongated in form (Fig. 9). In favourable cases one can see that the faintly yellowish drops or granules of which it now chiefly consists are becoming detached from the periphery of the main body of the elaioplast, and that these drops or granules are gradually scattered through the cell cavity (Fig. 10). Here they deepen their yellow tint, and in association with the red pigment developed in the cell-sap they produce the yellow, orange, or red coloration of the mature corolla-hairs, according as the one or the other pigmenting material predominates (Fig. 11).

The constituent plastids of these elaioplasts, therefore, undergo quite a similar series of changes as the chlorophyll grains in autumn leaves, which were first described in detail by Sachs in 1863 (10), or in ripening fruits, also studied by Sachs (1865). By the time the hairs are fully matured the elaioplast has entirely resolved itself into the scattered yellow pigment of the cell.

The corolla-hairs are not the only place in which elaioplasts occur in *Gaillardia*. They are also to be found in the stigmatic hairs, or in the more internal cells of the stigma and of the style, in the vegetative cells of the anther, and in the cells of the young pappus (calyx).

I sought for them in vain in the root-hairs or in the tissues of the root, in the leaf and the hairs which cover it, in the stem, and its clothing of hairs.

After I had completed my observations on the elaioplasts in the corolla-hairs, and drawn from them the conclusions which I have expressed above, I received a beautiful confirmation of the correctness of these views from the study of the elaioplasts in the other floral regions of *Gaillardia*.

In the cells of the connective of the young stamen, bodies occur which resemble the elaioplasts of the corolla-hairs in every respect except that they are coloured more or less deeply green. They are mostly spherical, although sometimes elongated in shape (Fig. 14). Moreover, in neighbouring cells of the connective we find every transition between deeply green bodies of this description, and others which are almost colourless and differ in no way from the elaioplasts of the corolla-hairs. On the addition of Iodine solution the occurrence of starch within the green bodies is readily demonstrated (Fig. 12). After remaining in the Iodine solution for some hours these intensely black-stained bodies form a most conspicuous feature in the otherwise yellow cells.

The appearance and reactions of these green bodies, no less than the transitions which occur between them and the ordinary elaioplasts, leave no doubt that they also are elaioplasts which contain chlorophyll, and which have retained the power of starch-formation. The cells of the young style and stigma also possess green, starch-producing elaioplasts.

Another very interesting case of green elaioplasts is furnished by the cells of the flattened basal plates or wings of the young pappus. In

many of these cells scattered chloroplasts occur, arranged as in ordinary assimilating tissue (Fig. 15). These chloroplasts are large, and many of them contain droplets of an apparently oily nature embedded within their substance. Probably these oil-drops mark the first stage of degeneration, although the power of starch-formation has not yet been lost. I was at first inclined to believe that the oily drops within these chlorophyll corpuscles represented the normal grana of these bodies developed to a rather unusual extent. The fact, however, that the chloroplasts of the other organs of *Gaillardia* (e.g., of the leaf) do not show any distinct grana of this kind, coupled with the further fact that the chlorophyll bodies of the pappus soon show undoubted signs of degeneration, has led me to conclude that the oil-drops are associated with the degradation of these chloroplasts.

In other cells of the pappus-plates the chlorophyll corpuscles tend to hang, more or less loosely, together. In yet other cells the aggregation of the chloroplasts is closer, although the outlines of each separate plastid is still maintained (Fig. 16). A further step in this aggregation of the chlorophyll bodies is seen in other, neighbouring cells in which they become so closely clumped together that the outlines of the individual chloroplasts can no longer be distinguished, and we obtain a typical green elaioplast in which the oil-drops of the plastids produce the finely granular appearance characteristic of these structures.

All these stages may be observed in adjoining cells of one and the same pappus-plate. They are best studied at about the time when the young pollen-grains are still without a membrane of their own and are enveloped in the special-wall (special mother-cell stage).

In older pappus-plates the green colour of the elaioplasts gives place to yellow, and other degeneration processes become evident.

Now that the development of the elaioplast has been followed in at least one species we are in a better position to compare this body with the oil-bodies of Hepaticae. Wakker evidently believed in the identity of the two structures, whilst other authors—such as Von Küster (11)—held an opposite opinion. On comparing what has been written above regarding the elaioplasts of *Gaillardia* with Garjeanne's careful account of the development of the oil-bodies of several Jungermanniales, it will be seen that the two structures have a very different origin. In the latter the oil-bodies arise as vacuoles in the cytoplasm, whilst we have seen that the elaioplasts of *Gaillardia* are formed by the aggregation of plastids and their degeneration products. Whilst, therefore, we cannot draw general conclusions until other species have been examined more fully, we may say that the developmental history of the elaioplasts of *Gaillardia* is essentially different from that of the oil-bodies of the Hepaticae.

External conditions seem to exert very little influence on the appearance of the elaioplasts.

I have kept the young capitula in total darkness for several (3-6) days without altering the development or structure of the elaioplasts of the corolla-hairs in the least.

The only deviation which I have ever found in the behaviour of the plastids of the corolla-hairs occurred in a very young capitulum from which the protecting bracts had been dissected away so that the tiny flower-buds were exposed to the full effect of the light.

Here the aggregation of the plastids into elaioplasts had been retarded in a number of cells.

The clumping together of the plastids of a cell into a more or less close mass is by no means an unusual occurrence. Kraus (12) many years ago described the effect of cold upon the chlorophyll-grains of winter leaves. Here these bodies were found to have passed from the walls to the interior of the cells and were there aggregated in clumps. Charles Darwin (13), in 1882, observed a very close massing of the chloroplasts in the cells of certain insectivorous plants under the influence of ammonium carbonate (a solution of 4-7 parts of ammonium carbonate in 1,000 parts water).

The work of Stahl (14), as well as of others, has shown that an irregular aggregation of chloroplasts is produced under the influence of intense illumination.

Pfeffer (15) mentions that similar results are induced by injuries and various mechanical agencies.

The close massing of the plastids into compact elaioplasts is most probably connected with their degeneration, and may very likely be compared to the aggregation of these bodies produced by the injurious agencies enumerated above. That the elaioplasts have any particular function to perform which is of direct significance to the life of the cell is most unlikely.

A secondary use for the degeneration products of the plastids—massed into elaioplasts—certainly does occur in the case of the corolla-hairs of *Gaillardia*, for here they give rise to the yellow pigment which forms an important part of the attractive apparatus of the mature flower.

In other situations, however, the elaioplasts seem to disappear, without having even this secondary biological significance.

It will be interesting to examine the Monocotyledonous elaioplasts again more closely in the light of what has been learnt of these bodies in *Gaillardia*, to see whether they possess the same nature and history. I hope to obtain material for this purpose during the next season.

In conclusion, I must express my indebtedness to a Government grant for assistance in carrying out this research.

SUMMARY.

1. Elaioplasts which hitherto had only been met with in Monocotyledons (and *Psilotum*) have now been found to occur in a Dicotyledon—*Gaillardia*.

2. The elaioplasts occurring in the corolla-hairs of *Gaillardia* are found to agree in their appearance and in their reactions with the elaioplasts described by Wakker and Zimmermann in Monocotyledons.

3. They have been found in the corolla-hairs, the pappus, the connective of the stamens, the style and the stigma of *Gaillardia*. They are absent from the tissues of the stem, the root, and the leaf of this plant.

4. They are formed by the aggregation of plastids and their degeneration products at one or more spots in the cell.

5. In the corolla-hairs of *Gaillardia* they give rise to the oily, yellow pigment which, in association with the red cell-sap, gives the mature hairs of the flower their characteristic colour.

6. The elaioplasts occurring in the stamens and in the style and stigma of *Gaillardia* agree in all respects with those of the corolla-hairs except that they are coloured green with chlorophyll, and can form starch within their substance. In neighbouring cells of these tissues all transitions occur between elaioplasts, which are coloured brightly green, and those which are almost colourless like those of the corolla-hairs.

7. In the tissues of the young pappus every transition can be found in neighbouring cells between those which contain scattered chloroplasts entirely free from one another, and those in which the chloroplasts have clumped together to form a green mass identical with the green elaioplast of the stamen or the stigma.

8. The elaioplasts of *Gaillardia* (and probably of the Monocotyledons also) differ essentially in their development from the oil-bodies of the liverworts.

9. External conditions were found to exert very little influence upon the appearance of the elaioplast, although rather strong, direct illumination seemed in one case to have somewhat retarded the aggregation of the plastids.

10. The close massing of the plastids into compact elaioplasts is probably connected with their degeneration, and may be compared to the aggregation of the plastids under the influence of various (mostly injurious) agencies described by several previous writers.

11. It is most unlikely that the elaioplasts perform any function of direct importance to the life of the plant, although they may in some cases (corolla-hairs of *Gaillardia*) serve a secondary, biological purpose.

REFERENCES TO LITERATURE.

1. WAKKER, J. H.: Studien über die Inhaltskörper der Pflanzenzelle. Pringsh. Jahrb. f. wiss. Bot., xix, 1888, p. 423, &c.
2. ZIMMERMANN, A.: Ueber die Elaioplasten. Beiträge z. Morphol. u. Physiol. d. Pflanzenzelle, 1893, p. 185, &c.
3. RACIBORSKI, M.: Ueber d. Entwicklungsgesch. d. Elaioplasten d. Liliaceen. Anzeig. d. Akad. d. Wiss. in Krakau, 1893, p. 259. I was unable to obtain this paper and quote from the abstract in Bot. Centralbl., lvii, 1894, p. 305.
4. PFEFFER, W.: Die Oelkörper der Lebermoose. Flora, 1874, p. 2, &c.
5. GARJEANNE, M. J.: Die Oelkörper der Jungermanniales. Flora, xcii, 1903, p. 457, &c.
6. RADIKOFER, L.: Zur Klärung von Theophrastus und der Theophrasteen. Sitzungsber. d. math.-physik. Kl. d. K. B. Akad. d. Wiss. zu München, xix, 1889, p. 221.
7. ———: Ueber die Gliederung d. Familie d. Sapindaceen. Ib., xx, 1890, p. 105.
8. MONTEVERDE, N. A.: Ueber die Ablagerung von Calcium- und Magnesium-Oxalate in d. Pflanze. Petersburg, 1889.
9. SOLEREDER, H.: Studien über die Tribus der Gaertnereen. Ber. d. Deutsch. Bot. Gesellsch., 1890, p. 71.
10. SACHS, J. VON: Entleerung der Blätter im Herbst. Flora, 1863, p. 200.
11. KÜSTER, VON: Die Oelkörper der Lebermoose u. ihr Verhältniss zu den Elaioplasten. Basel, 1894.
12. KRAUS, G.: Bot. Zeit., 1874, p. 406.
13. DARWIN, C.: The action of Carbonate of Ammonia on Chlorophyll Bodies. Journ. Linn. Soc., xix, 1882, p. 262.
14. STAHL, E.: Ueber den Einfluss d. Richtung u. Stärke d. Beleuchtung auf einige Bewegungserscheinungen im Pflanzenreiche. Bot. Zeit., 1880, p. 340.
15. PFEFFER, W.: Physiology of Plants (Eng. trans.), iii, 1906, p. 379.

EXPLANATION OF FIGURES IN PLATE IV.

Illustrating Mr. Beer's Paper on Elaioplasts.

All the figures refer to tissues of *Gaillardia Lorensiana*. Preparations examined and drawn in .6 per cent. NaCl solution unless otherwise stated.

Fig. 1. Very young cell of corolla-hair before elaioplasts have developed. Leucoplasts distributed in the cytoplasm. $\times 1075$.

Fig. 2. Apex of another corolla-hair showing scattered leucoplasts. $\times 650$.

Fig. 3. Young cell of corolla-hair showing an early stage in the aggregation of plastids. $\times 650$.

Fig. 4. Corolla-hair showing a stage in aggregation of plastids. $\times 650$.

Fig. 5. Corolla-hair with later stage of elaioplast development. $\times 650$.

Figs. 6, 7, 8. Elaioplast completely developed in corolla-hairs. $\times 650$.

Fig. 9. Cell from older corolla-hair. Elaioplast elongated and showing first indication of disintegrating. $\times 650$.


Fig. 10. Later stage; substance of elaioplast becoming distributed through the cell. $\times 600$.

Fig. 11. Mature corolla-hair. Yellow pigment-material derived from elaioplast completely distributed through the cell. $\times 650$.

Fig. 12. Green elaioplast from anther after two days in dilute Iodine solution. Oil-drops have been exuded at the surface. Starch (shown in the figure as black grains) contained in the yellow-stained matrix. $\times 600$.

Fig. 13. Elaioplast from corolla-hair in .6 per cent. NaCl after gently warming on slide over spirit-lamp. $\times 650$.

Figs. 15 and 16. Two neighbouring cells from plate of pappus. In Fig. 15 the chloroplasts are scattered, whilst in Fig. 16 they are massed into a loose clump. Note oil-drops within the chloroplasts (represented as black dots). $\times$ about 600.



"On Elaioplasts"

B7a

By Rudolf Beer BSc. F.L.S.

In 1888 Wakker⁽¹⁾ described the occurrence of a conspicuous body lying in the epidermal cells of the leaves + in the superficial tissues of the root + stem of Vanilla planifolia. This body is somewhat larger than the nucleus + considerably larger than the amyloplasts; it possesses a ^{sharply} ~~well~~ defined outline + a peculiar, ^{somewhat} ~~absent~~ yellowish colour. In each plate-like epidermal cell one such body occurs + it often lies near the nucleus although ^{in other cases} it may occupy a different position in the cell. As this body consists of protoplasmic material + contains oil Wakker ~~proposed to name it~~ elaioplast or oil-former. ~~He also found elaioplasts to occur in another species of Vanilla viz one known~~
~~as Vanilla aromatica latifolia He~~
He made a careful study of the effect of reagents upon the elaioplast ^{briefly} + describes their gradual disappearance ^{of these bodies} in older cells. Thus

2
~~Thus on 8th January he found the epidermal cells of a growing leaf to contain elaioplasts measuring 12 μ in diameter. Three weeks later they only measured 6-8 μ ; ~~which~~ after a further three weeks interval only a few cells ^{were found to} contained elaioplasts at all whilst in the majority only scattered oil drops occurred in the cytoplasm.~~
~~With regard to the origin of the elaioplasts in the young cells~~ Wappler was unable to study the origin of the elaioplasts ~~on account of~~ as he had no suitable material for this purpose.

In 1893 ~~Zimmerman~~ ^{found} similar ^{also} bodies to occur in the ~~also~~ found elaioplasts ^{to occur} in the cells of another species of *Vanilla*, ~~which was known to him under the name of~~ *Vanilla aromatica latifolia*.
In 1893 Zimmerman ⁽²⁾ found similar bodies to occur in *Funkia coerulescens*, *F. lancifolia*, *F. Sieboldiana*, *Dracaena* sp., *Ornithogalum scilloides*, *Agave americana*, *A. mitis* + in *Oncidium suave*. // Raciborski ⁽³⁾ in the same year described elaioplasts in the tissues of various species of *Ornithogalum*, *Albuca*, *Funkia* + *Gagea*.
Zimmerman further found these bodies

~~masses~~ in the internal cells of the stem of Psilotum
 & in the perianth leaves of Maschallaria picta.

The shape of the elaioplasts ^{differs} ~~is different~~ in the
 various plants but it is usually constant in the
 same species. Spherical forms, grape-like
 bodies, irregular plasmodium-like masses
 have all been described.

Usually only one elaioplast occurs in a cell
 but in some cases they may be more numerous
 (e.g. Ornithogalum). ~~In structure~~ The finer

structure of the elaioplast has been ^{carefully} examined both by Zimmermann & Raiborski.

It ~~possesses~~ ^{has} a finely granular appearance
~~which is~~ due to the occurrence of a number
 of tiny, highly refractive spherical bodies
 lying in its substance. One ~~one~~ or more
 less refractive spots often occur within the
 elaioplast. The elaioplasts of the species of
Gagea are said by Raiborski to possess
 a somewhat different structure. ~~These~~
 In these plants the usually spherical elaioplasts are

characterized by the feeble development of the stroma,
~~consist of spherical bodies~~ ^{they are} surrounded by
 a plasmatic envelope ^{& contain within this} enclosing an oily
 substance like that found in other elaioplasts.

The micro-chemical observations of Wacker &
 Zimmermann have shown that the elaioplast
 consists of a plasmatic matrix or stroma in
 in which are embedded a number of numerous
 minute oil drops. Zimmermann points out
 that these oil drops show a close similarity in
 their reactions with the oil drops obtained from
 plastids.

The origin of the elaioplasts has been very
 little studied. Raiboroki states that in
Ornithogalum umbellatum they arise as small
 highly refractive spherules which always lie
 at one pole of the usually elongated
 cell-nucleus.

Although it is known that in older cells the
 elaioplasts disappear their fate has never

Although Wakker had no material with which to
work out the development of the elaioplast. He
ventures an ~~inadequate~~ interesting suggestion at the
conclusion of his account of the oil-bodies of
liverworts "Leider ist durch diese Mittheilung
"der Ursprung der Elaioplasten nicht ausgemacht,
"es ist mir aber äusserst wahrscheinlich, dass
"es bei den Lebermoosen metamorphosirte
"Chlorophyllkörner sind. Vielleicht ist dieses
"auch bei Vanilla der Fall".

Digitized by Hunt Institute for Botanical Documentation
~~As Garjeanne showed the suggestion does~~
~~will not apply to the oil-bodies of the liverworts~~
~~but~~ Garjeanne's work (5) on the oil-bodies of
the Jungermanniales ~~proves~~ ^{have} clearly indicated
that these bodies ~~had~~ ^{have} a different origin to that
suggested by Wakker; my own observations
of on elaioplasts recorded below will, ^{however,} show that
his guess was ^{nearer the truth,} ~~happens~~ in the case of some
for Phanerogamic elaioplasts.

6
has definitely traced.

The function of the chloroplast is quite unknown. Wacker believed ^{that they might} ~~they~~ be oil-formers much as plastids leucoplasts are starch-formers. Zimmerman ~~once~~ offered the suggestion that they might ^{possibly} be parasitic or symbiotic fungi living within the cells of the higher plant.

Raiboraki, however, considers them to be normal organs of the cell which contains them + classes them with oil-bodies, tannin-vesicles + ordinary vacuoles.

In the liverworts oil-bodies have been known to occur since the time of Gottsche + even earlier. ~~They were first described in detail by Halle~~ The first really fundamental description of these bodies is due to Pfeffer⁽⁴⁾ + quite recently their development has been fully worked out by Garjeanne⁽⁵⁾. ~~Opinion has been~~ Oil or fat bodies of a somewhat similar

7

to those of the liverworts
appearance have been described by Radlkofer (6)
Monteverde (8), Solereder (9) + others in the tissues of
a number of the Dicotyledons + Monocotyledons.
Opinion ~~has varied~~ ^{varies} very much as to the
relation of these oil-bodies of liverworts +
Phanerogams to the elaioplasts. Some believe
the two structures to be ^{closely allied} whilst others are
of opinion that they are ^{radically} ~~of quite a~~ different
~~nature~~ from one another.

On p. 458 of ^{Gargeanne's} ~~his~~ work on the oil bodies of
Fingermanniales ~~Gargeanne~~ ^{Gargeanne} there
occurs a sentence which is of considerable
interest in the light of what is to follow in the
present paper. He ~~says~~ in referring to Wacker's
elaioplasts he says "These oil-formers or
Elaioplasts are generally to be found in the
neighbourhood of the nucleus & agree in ~~idea~~ in
their young state agree in several respects with
leucoplasts". He enters into no details, however,
^{does not give his} ~~gives no~~ reasons for this assertion.

From this brief survey of the literature it will be seen that elaioplasts corresponding to those discovered by Wacker in Vanilla have up to the present been found only in Monocotyledons with the single exception of Psilotum reported by Zimmermann.

Moreover, apart from a few quite insufficient observations we are still entirely in the dark regarding their true nature, their significance, their origin & their fate.

During my observations upon the pollen grains of Compositae I was interested to find bodies which resembled the Monocotyledonous elaioplasts occurring in the tissues of the floral region of Saillardia ^(a) ~~in~~ Lorenziana. ~~of the garden~~

This would extend the distribution of elaioplasts to a member of the Dicotyledons.

My first endeavour was to make

-
- (a) Saillardia Lorenziana is a German variety of G. picta which itself appears to be only a garden ~~for~~ variety of G. pulchella.

I have not yet had an opportunity of examining any other variety form or species of Saillardia except the one mentioned above.

I was dealing with
 sure that ¹ true elaioplasts. In the
 hairs upon the perianth of young flowers, where
 I first observed these bodies, they occur as
 more or less spherical, highly refractive, ~~granular~~
 granular structures usually somewhat larger than
 the nucleus & in most cases lying singly in a
 cell although two or even more such bodies were
 occasionally met with.

Heated gently upon the slide (whilst lying in a
 physiological salt solution) drops of oil were
 exuded from their surface; in osmic acid
 (as this occurs in Flemming's stronger solution) they
 are blackened; alkanin solution is 60%
 alcohol colours the bodies deeply red; Iodine
 solution (in KI) colours them brown; absolute
 alcohol dissolves out the oil from their interior
 & leaves them vacuolated; in nitric acid (warmed)
 + subsequent addition of ammonia they give the
 xanthoproteic reaction;

Their reactions were found to be as follows:—

- (1) Heated gently upon the slide (whilst lying in a physiological salt solution) drops of oil are exuded from their surface (Fig 13)
- (2) Osmic acid (as this occurs in Flemin's stronger solution) turns them black or brown.
- (3) Potassium bichromate after 24 hours action leaves them quite colourless but causes the extrusion of oil globules [as in (1)].
- (4) Alkannin solution in 60% alcohol colours the bodies deeply red.
- (5) Iodine (in K.I.) colours them brown & causes the extrusion of oil-globules (Fig 12)
- (6) Absolute alcohol washes out the oil from their interior & leaves them vacuolated.
- (7) 2 strong HNO_3 (warmed) followed by NH_3 they give the xanthoproteic reaction (viz yellow colouration).
- (8) Glacial acetic acid after 24 hours action causes great extrusion of oil drops but no solution.

(9) 10% KOH. after 24 hours action dissolves
neither stroma nor oil drops.

These reactions, combined with their general appearance,
~~will~~ show that the bodies occurring in the ^{hairs} ~~cells~~
of *Gaillardia* are in all respects similar to
the leucoplasts described by previous authors.

In very young hairs from capitula which were
still quite small & immature & entirely ^{enclosed within} ~~covered~~
~~the~~ ~~insoluble~~ ~~tracts~~ no leucoplasts were
yet to be seen. The cell contained a nucleus
& ~~vacuolated~~ cytoplasm which partly formed a
peripheral layer & partly extended ~~as in~~ ^{through the cell cavity.} in
strands & bars ~~to the nucleus~~ (Figs 1 & 2)

Embedded in the cytoplasm were a number of
very small highly refractive grains which had
all ^{the} ~~very~~ appearance of ordinary leucoplasts.

~~The fact that~~ That these refractive grains
are really leucoplasts is ^{confirmed} ~~shown~~ by two facts.

12
Firstly the resemblance between the unquestionable
starch-forming leucoplasts occurring ^{for example,} in the
hairs ^{which cover} the very young leaves & the highly
refractive grains ^{contained} in the perianth-hairs is complete
although starch is not found ^{in the latter} under the ~~the~~ usual
conditions ~~in the latter~~ of growth.

Secondly if the enveloping bracts be removed from
a ~~very~~ young flower ~~and~~ ^(without detaching it from the parent plant) the
hairs upon the perianth-hairs exposed to
a strong insolation starch can be seen
to have developed in ~~some~~ ^{some} of these
refractive ~~granules~~ grains.

For these reasons I believe the highly refractive
grains occurring in the cells of the perianth-hairs
to be leucoplasts, ~~or products~~ some of which,
however, may have lost the ~~sample~~ power
of starch formation.

In somewhat older hairs these plastids, ^{a number} ~~some~~ of which show signs of undergoing degeneration, tend to aggregate together at one or more spots within the cell. ~~Some~~ Not infrequently the aggregation of the plastids is in the neighbourhood of the nucleus but in many instances it is found to occur at other regions of the cell. (Figs 3 + 4)

At first the aggregation of the refractive grains is a very loose one but it gradually grows closer & closer ^(Fig 5) until ~~bodies~~ compact, highly refractive bodies are formed, which we have already recognised as Elaisoplasts, ^(Figs 6, 7 & 8). The elaisoplasts in the penultimate hairs of Gaillardia are, therefore, formed by the aggregation of plastids & their ^{degeneration} products, ~~degeneration~~ ^{then being} ~~possibly associated with some~~ cytoplasm as well, at one or more spots in the cell.

Within the elaisoplast ~~I believe~~ the plastids appear soon to undergo further degeneration with the

production of an oily material. That
Timmermann should find a close similarity
between the oil of the elaioplasts studied by him
+ the oil obtained from plastids is no
longer surprising.

All the plastids ~~the~~ of the cell have not
clumped together within the elaioplast. a
certain proportion ~~are~~ ^{remains cultured} ~~are~~ ^{generally} ~~still~~ ⁱⁿ ~~the periphery~~
through the cell. (Fig 7).

~~It seems probable that these plastids
which are beginning to degenerate become
massed together in this manner whilst
others which~~

~~For some time there is little alteration within
the cell. In considerably older hairs,
however, we find~~

For some time there is little ~~old~~ alteration within
the cell. The conspicuous elaioplast may
lie in almost any part of the cell but
often it takes up a position near the

nucleus. In some instances it entirely envelopes the nucleus as I have represented in Fig 8.

In much older hairs we find the elaioplast undergoing a change. Its outline becomes less regular & in some cases it becomes drawn out & elongated in form (Fig 9).

In favourable cases one can see that ~~the~~ ^{the drops or granules} faintly yellow ~~is~~ ^{of which it now chiefly} ~~it~~ ^{periphery of the} ~~it~~ ^{main} ~~it~~ ^{body} ~~it~~ ^{of the elaioplast} ~~it~~ ^{these drops or granules} ~~it~~ ^{are gradually becoming} ~~it~~ ^{scattered} ~~it~~ ^{through the cell cavity} ~~it~~ ^(Fig 10)

are gradually ~~scattered~~ ^{scattered} through the cell cavity (Fig 10). Here they deepen their yellow tint & in association with the red pigment developed in the cell-sap they produce the yellow, orange or red coloration of the mature perianth-hairs according as the one or the other pigmenting material predominates. (Fig 11).

The constituent plastids of these elaioplasts, therefore, undergo ~~a~~ quite a similar series of changes as the chlorophyll grains in autumn leaves ^{which were} first described in detail by Sachs in 1863 (10) or in ripening fruits also studied by Sachs (1865) ~~as well as others~~.
~~After leaving~~ After By the time the hairs are fully matured ~~the~~ the elaioplast has entirely resolved itself into the scattered yellow pigment of the cell.

Digitized by Hunt Institute for Botanical Documentation

The perianth-hairs are not the only place in which elaioplasts occur in *Saillardinia*. ~~In the stigmatic hairs~~ They ^{also} are to be found in the stigmatic hairs, in the more internal cells of the stigma & of the style, in the vegetative cells of the anther & in the cells of the young pappus (calyx).

I sought for them in vain in the root-hairs of ⁱⁿ the tissues of the root, in the leaf or

17

+ the hairs which cover it, in the stem & its clothing of hairs.

After I had completed ^{my observations on} ~~the study of~~ the elaioplasts in the perianth-hairs ~~of~~ & drawn ^{from them} the conclusions which I have expressed above I received a beautiful confirmation of the correctness of ~~my~~ ^{these} views from the study of the elaioplasts in the other floral regions of Gaillardia.

In the cells of the connective of the ^{young} stamen bodies occur which resemble the elaioplasts of the perianth-hairs in every respect except that they are colored more or less deeply green. They are mostly spherical although sometimes elongated in shape. (Fig 14)

Moreover in neighboring cells of the connective we find every transition between ~~these~~ deeply green bodies of this description & others which are ^{almost} ~~quite~~ colorless & differ in

no ^{way} respect from the elaioplasts already of the perianth-hairs. On the addition of Iodine solution the occurrence of starch within the green bodies is readily demonstrated. ~~the less the green pigment~~ (Fig 12).

After remaining in the Iodine solution for some hours these ^{intensely} ~~deeply~~ black stained bodies form ^a the most conspicuous feature in the otherwise yellow cells.

The appearance + reactions of these green bodies, no less than the transitions which occur between them + the ordinary elaioplasts, leave no doubt that they ^{also} are elaioplasts which ^{contain} ~~have developed for~~ ~~retained~~ chlorophyll + ^{which have} retained the power of starch formation. The cells of the ^{young} style + stigma also possess green, starch-producing elaioplasts.

Another very interesting ~~position~~ ^{case of} green elaioplasts is furnished by ^{cells of the} ~~the~~ flattened basal plates ^{or wings} of the young pappus.

In many of these cells scattered chloroplasts occur arranged as in ordinary assimilating tissue ^(Fig 15). These chloroplasts are large & ^{many of them} contain ~~drops~~ droplets of an apparently oily nature ~~within~~ embedded within their substance. ^{these} Probably these oil-drops ~~are~~ mark the first stage of degeneration although the ~~assimilative~~ power of starch-formation has not yet been lost. I was at first inclined to believe that the oily drops within these chlorophyll corpuscles represented the normal grana of these bodies developed to a rather unusual extent. The fact, however, that the chlorophyll chloroplasts of the other organs of *Gaillardia* (e.g. of the leaf) do not show any distinct grana of this kind coupled with the ~~fact~~ further fact that the ^{chlorophyll bodies} ~~plastids~~ of the pappus soon ~~enough~~ show undoubted signs of degeneration has lead me to conclude that the oil-drops are associated with the degradation of these chloroplasts.

In other cells lying quite near to those with

Scattered chloroplasts are others in which these bodies tend to hang, more or less loosely, together. In other cells again the aggregation of the chloroplasts is closer although the outlines of each separate plastid is still maintained ^(Fig. 16). A further step in the aggregation of the chloroplast bodies is seen in other neighbouring cells in which they become so closely clumped together that the outlines of the individual chloroplasts can no longer be distinguished & we obtain a typical green chloroplast in which the oil drops of the plastids produce the finely granular appearance characteristic of these structures.

All these stages may be observed in adjoining cells of one & the same pappus-plate. They are best studied at about the time when the young pollen grains are still without a membrane of their own & are enveloped in the special-wall (special mother-cell stage)

In older pappus-plates the green colour of the elaioplasts gives place to yellow & other degeneration processes become evident.

Now that the development of the elaioplast has been followed in at least one species we are in a better position to compare this body with the oil-bodies of Hepaticae. Wacker evidently believed in the identity of the two structures whilst other authors - such as Von Klüber⁽¹¹⁾ ~~has nothing~~ held an ^{opposite} ~~different~~ opinion.

On comparing what has been ~~not~~ written above ~~with~~ regarding the elaioplasts of Gaillardia with Larijé's careful account of the development of the oil-bodies of several Jungermanniales it will be seen that the two structures have a very different origin. In the latter the oil-bodies arise as vacuoles in the cytoplasm whilst we have seen that the elaioplasts of

Gaillardia are formed by the aggregation of plastids & their degeneration products. Whilst, therefore, we cannot ~~as yet~~ draw general conclusions as ^{until other species have been examined more fully} ~~to all elaioplasts~~ we may say that the developmental history of the elaioplasts of Gaillardia is essentially different from that of the oil bodies of the Hepaticae.

External conditions ^{seem to} ~~not~~ exert very little influence on the appearance of the elaioplasts.

I have kept the young ~~flower buds~~ capitula in total darkness for ^{several (3-6)} ~~for three to six~~ days without altering the ~~structure~~ ^{development} or structure of the elaioplasts of the persistent hairs in the leaf.

The only deviation which I have ever found in the behaviour of the plastids of the persistent hairs occurred in a very young capitulum ~~about~~ from which the protecting ~~buds~~ bracts had been dissected away so that the ^{young} flower-buds were

exposed to the full effect of the light.

Here the aggregation of the plastids into
elaioplasts had been retarded in a number of
cells.

The clumping together of the plastids of a cell into
a more or less close mass is by no means an
unusual occurrence. Kraus (12) many

years ago described the effect of cold upon the
chlorophyll-grains of winter leaves. Here these
bodies were found to have passed from the walls to the
interior of the cells & were there aggregated in clumps.

Charles Darwin (13) in 1882 observed a very
close massing of the chloroplasts in the cells
of certain insectivorous plants under the
influence of ammonium carbonate (a solution
of 4-7 ^{parts of} ammonium carbonate ⁱⁿ the 1000 ^{parts} water)

The work of Stahl, ^{as} as well as of others, has shown that an irregular aggregation of the chloroplasts is produced under the influence of intense illumination.

Dr. Pfeffer ⁽¹⁵⁾ mentions that similar results are induced by injuries & various mechanical agencies.

The close massing of the plastids into compact chloroplasts is most probably connected with their degeneration & may ^{very} ~~not~~ likely be compared to the aggregation of these bodies produced by the injurious agencies enumerated above. That

the chloroplasts have any particular ~~useful~~ part to play in the function to perform which is of direct significance to the life of the cell is most unlikely. ~~They are structures which~~

~~The plastids are no longer useful required of importance to the cell as assimilating or amylloplastic organs & the~~

A secondary use for the degeneration products of the plastids - moved into chloroplasts - ~~may~~ is certainly does occur in the case of the perianth-

hairs of Gaillardia for here they give rise to
 the yellow pigment which ^{is important} forms an ^{important} part of the
 attractive apparatus of the mature flower.
 In other situations, ^{however} the elaioplasts ~~app~~ ^{seem to} disappear
~~however~~ without having even this ^{secondary} ~~chance~~
 biological significance.

It will be interesting to examine the Monocotyledonous
 elaioplasts again more closely in the light of
~~the above~~ what has been learnt of these bodies
 in Gaillardia to see whether they ~~are~~ ^{possess} the
 same nature & history. I hope to obtain
 material for this purpose during the next
 season.

In conclusion I must express my
 indebtedness to a ~~the~~ Government Grant ~~for~~
~~for a Term 2-4 to N.A. oil immersion~~
~~lens which has greatly facilitated my~~
~~observations.~~
 for assistance in carrying out this research.

26

Summary

- (1) Elaioplasts which hitherto had only been met with in Monocotyledons (+ Poilatum) have now been found to occur in a Dicotyledon - Gaillardia -.
- (2) The elaioplasts occurring in the perianth-hairs of Gaillardia are found to agree in ~~their~~^{their reactions} their appearance & in ~~their~~ behaviour towards reagents with the elaioplasts described by Wakker & Zimmermann in Monocotyledons.
- (4) They have been found in the perianth-hairs, the pappus (~~in an early stage of development~~), the connective of the Stamens, the style & the stigma of Gaillardia. They are absent from the tissues of the stem, the root & the leaf of this plant.
- (5) They are formed by the aggregation of ~~the~~ ~~elaioplasts~~ plastids of the cell & their degeneration products at one or more spots in the cell.
- (6) In the perianth-hairs of Gaillardia they give

27

rise to the oily, yellow pigment which in association with the red cell-saps gives the mature hairs of the flower their characteristic colour.

- (7) The elaioplasts occurring in the stamens & in the style & stigma of Gaillardia agree in all respects with those of the perianth-hairs except that they are coloured green ^{with chlorophyll} (can) form starch within their substance. In ~~these~~

neighbouring cells of these tissues all transitions occur between elaioplasts which are coloured brightly green & those which are almost colourless like those of the perianth-hairs.

- (8) In the tissues of the ~~plate of the~~ young pappus every transition can be found in neighbouring cells between those which contain scattered chloroplasts entirely free from one another & those in which the chloroplasts have clumped together to form a ~~mass~~ green mass identical with the green elaioplasts of the stamen or the stigma.

(9) The elaioplasts of Gaillardia (+ probably ^{of the} ~~other~~ ^{other} ~~Monocotyledonous~~ ~~elaioplasts~~ also) differ essentially in their development from the oil-bodies of the biennials.

(10) External conditions were found to exert very little influence upon the appearance of the elaioplast although ^{rather} ~~intense~~ strong direct illumination seemed in one case to have ^{somewhat} retarded the aggregation of the plastids.

(11) The close massing of the plastids into compact elaioplasts is probably connected with their degeneration & may be compared to the aggregation of the plastids under the influence of various ^(mostly) ~~injurious~~ agencies described by ^{several previous} ~~several~~ writers.

(12) It is most unlikely that the elaioplasts ^{perform} ~~have~~ any function of direct importance to the life of the plant although they may ~~be~~ in some cases (perianth-hairs of Gaillardia) & serve a secondary, biological purpose.

References to Literature.

- (1) Wakker J.H. "Studien über die Inhaltskörper der Pflanzenzelle" *Pringsh. Jahrb. f. wiss. Bot.* Bd 19 1888 p. 423 etc.
- (2) Zimmermann A. "Ueber die Elaioplasten" *Beiträge z. Morphol. + Physiol. d. Pflanzenzelle* 1893 pp. 185 etc.
- (3) Raichevski M. "Ueber d. Entwicklungsgesch. d. Elaioplasten d. *Elaiaceen*" *Anzeig. d. Akad. d. Wiss. in Krakau* 1893 p. 259
 I was unable to obtain this paper & quote from the abstract - *Bot. Centrall. Bd LVII* 1894 p. 305
- (4) Pfeffer W. Die Ölkörper der Lebermoose *Flora* 1894 p. 2 etc.
- (5) Garjeanne M.J. "Die Ölkörper der Jungermanniales" *Flora* Bd 92 1903 p. 457 etc.
- (6) Radtkofer L. "Zur Klärung von Thesphrasta und der Thesphrasteen" *Sitzungsber. d. math.-physik. Kl. d. K. B. Akad. d. Wiss. z. München* Bd 19. 1889 p. 221
- (7) Radtkofer L. "Ueber die Gliederung d. Familie d. Sapindaceen" *Abh.* Bd 20 1890 p. 105

- (8) Monteverde N.A. "Über die Ablagerung von Calcium-
und Magnesium-Oxalate in d. Pflanze
Petersburg 1889
- (9) Solereder H. "Studien über die Tribus der
Gaertnereen." Ber. d. Deutsch. bot.
Gesellsch. 1890 p. 71
- (10) Sachs J. von. "Entleerung der Blätter im Herbst"
Flora 1863 p. 200
- (11) Küster von "Die Ölkörper der Lebermoose
u. ihr Verhältniss zu den Elaioplasten"
Basel 1894
- (12) Kraus G. Bot. Zeit. 1874 p. 406
- ~~73~~
- (13) Darwin C.
Journ. Linn. Soc. Vol 19 1882 p. 262
- (14) Stahl E. "Ueber den Einfluss d. Richtung u.
Stärke d. Beleuchtung auf einige
Bewegungsersch. im Pflanzenreiche"
Bot. Zeit. 1880 p. 340
- (15) Pfeffer W. Physiology of Plants (Engl. trans)
Vol 3 . 1906 p. 329

Explanation of Figures in Plate —

All the figures refer to tissues of Gaillardia
Lorentziana. ~~Such~~ Preparations examined & drawn
in .6% NaCl solution unless otherwise stated.

- Fig 1. Very young cell of perianth-hair before
elaioplasts have developed. Leucoplasts
distributed in the cytoplasm $\times 1075$
- Fig 2. Apex of another perianth-hair showing
scattered leucoplasts $\times 650$
- Fig 3. Young cell of perianth-hair showing an
early stage in the aggregation of plastids.
- Fig 4. Perianth-hair showing a ~~small~~ X 650
stage in aggregation of plastids $\times 650$
- Fig 5. Perianth-hair with later ~~stage~~ stage of
elaioplast development. $\times 650$
- Fig 6, 7, 8. Elaioplast completely developed
in perianth-hairs. $\times 650$
- Fig 9. ~~Older~~ Cell from older perianth-hair.
+ Elaioplast elongated + showing first

indication of disintegrating x 650

Fig 10 Later stage; substance of elaioplast becoming distributed through the cell.
x 600

Fig 11 Mature perianth-hair. Yellow pigment-material derived from elaioplast completely distributed through the cell.
x 650

Fig 12 Green elaioplast from anther after 2 days in dilute Iodine solution. Oil-drops have been exuded at the surface. Starch (shown in the figure as black grains) contained in the yellow stained matrix. x 600

Fig 13 Elaioplast from perianth hair in .6% NaCl after gently warming on slide over spirit-lamp x 650.

Fig 14 Green elaioplast in cell of anther
x 650

Figs 15 & 16. Two neighbouring cells from plate of pappus. In fig 15 the chloroplasts are scattered whilst in fig 16 they are ~~loosely~~ massed into a loose clump. Note oil drops

within the chloroplasts (represented as black dots)
X about 600.